arXiv:1202.1510v1 [math.PR] 7 Feb 2012

Spectral gap estimates at low temperature by decomposition of the energy landscape

Georg Menz, André Schlichting*

February 8, 2012

We consider a diffusion on a potential landscape which is given by a smooth Hamiltonian $H : \mathbb{R}^n \to \mathbb{R}$ in the regime of small noise ε. We give a new proof of the Eyring-Kramers formula for the spectral gap of the associated generator $L = \varepsilon\Delta - \nabla H \cdot \nabla$ of the diffusion. The proof is based on a refinement of the two-scale approach introduced by Grunewald, Otto, Westdickenberg, and Villani [GOVW09] and of the mean-difference estimate introduced by Chafaï and Malrieu [CM10]. The Eyring-Kramers formula follows as a simple corollary from two main ingredients: The first one shows that the spectral gap of the diffusion restricted to a basin of attraction of a local minimum scales nicely in ε. This mimics the fast convergence of the diffusion to metastable states. The second ingredient is the estimation of a mean-difference by a new weighted transport distance. It contains the main contribution of the spectral gap of L, resulting from exponential long waiting times of jumps between metastable states of the diffusion.

MSC: Primary 58J65; secondary 60J45; 60J60; 35P15; 49Q20.

Keywords: Eyring-Kramers formula, diffusion process, Kramers law, metastability, Poincaré inequality, spectral gap, weighted transport distance.

1. Introduction and main results

We consider a diffusion on a potential landscape which is given by a smooth Hamiltonian $H : \mathbb{R}^n \to \mathbb{R}$. We are interested in the regime of small noise ε. The generator of the

---

*Max Planck Institute for Mathematics in the Sciences, Leipzig, Germany.
Georg.Menz@mis.mpg.de, Andre.Schlichting@mis.mpg.de.

diffusion has the following form

$$L = \varepsilon \Delta - \nabla H \cdot \nabla. \tag{1.1}$$

The associated Dirichlet form is given by

$$\mathcal{E}(f) := \int (-Lf) f \,\mathrm{d}\mu = \varepsilon \int |\nabla f|^2 \,\mathrm{d}\mu. \tag{1.2}$$

The corresponding diffusion ξ_t satisfies the stochastic differential equation

$$\mathrm{d}\xi_t = -\nabla H(\xi_t)\ \mathrm{d}t + \sqrt{2\varepsilon}\ \mathrm{d}B_t, \tag{1.3}$$

where B_t stands for the Brownian motion on $\mathbb{R}^n$. The last equation is often also called *over-damped Langevin equation.* Under some growth assumption on H (cf. Section 1.1) there exists an equilibrium measure of the according stochastic process, which is called *Gibbs measure* and is given by

$$\mu(\mathrm{d}x) = \frac{1}{Z_\mu} \exp\left(-\frac{H(x)}{\varepsilon}\right) \mathrm{d}x \qquad \text{with} \qquad Z_\mu = \int \exp\left(-\frac{H(x)}{\varepsilon}\right) \mathrm{d}x. \tag{1.4}$$

We are particularly interested in the case where H has several local minima. Then, the process shows metastable behavior in the sense that there exists a separation of scales. On the fast scale, the process converges quickly to a neighborhood of a local minima. On the slow scale, the process stays there for an exponentially long waiting time after which it jumps to another local minima.

This behavior is well known in the context of chemical reactions. The exponential waiting time follows the *Arrhenius law* [Arr89] meaning that the mean exit time from one local minima of H to another one is exponentially large in the energy barrier between them. By now, the Arrhenius law is well-understood even for more complicated systems by the Freidlin-Wentzell theory [FW98], which is based on large deviations.

A refinement of the Arrhenius law is the *Kramers law* which additionally considers pre-exponential factors. The Kramers law goes back to Eyring [Eyr35] and Kramers [Kra40], which considered the one-dimensional situation. Both argued that also in high-dimensional problems of chemical reactions most reactions are nearby a single trajectory called *reaction pathway.* Evaluating the Hamiltonian along this *reaction coordinate* gives the classical picture of a double well potential in one dimension with an *energy barrier* separating the two local minima for which explicit calculations are feasible.

A rigorous proof of the Kramers law for the multidimensional case was open for a long time until Bovier, Eckhoff, Gayrard, and Klein [BEGK04, BGK05] obtained first order asymptotics that are sharp in the parameter ε. They also clarified the close connection between mean exit times, capacities and the exponentially small eigenvalues of the operator L given by (1.1). To set the nomenclature, for us the Eyring-Kramers formula is a sharp asymptotics for the exponentially small eigenvalues. The main tool of [BEGK04, BGK05] is potential theory. The small eigenvalues are related to the mean exit times of appropriate subsets of the state space. Further, the mean exit times are given by Newtonian capacities which can explicitly be calculated in the regime of small noise ε.

A little bit later Helffer, Klein, and Nier [HKN04, HN05, HN06] also deduced the Eyring-Kramers formula using the connection of the spectral gap estimate of the Fokker-Planck operator L given by (1.1) to the one of the Witten Laplacian. This approach makes it possible to get quantitative results with the help of semiclassical analysis. They deduced sharp asymptotics of the exponentially small eigenvalues of L and gave an explicit expansion in ε to theoretically any order. For a nice overview on the Kramer's law we refer the reader to the article [Ber11].

The aim of this article is to provide a new self-contained proof of the Eyring-Kramers formula for the first eigenvalue. Functional inequalities are the background of our approach. Our proof is based on two-basic ingredients:

- The first one is a refinement of the two-scale approach of Grunewald, Otto, Westdickenberg, and Villani [GOVW09] leading to a good Poincaré inequality for Gibbs measures μ restricted to a basin of attraction of a local minima (cf. Theorem 1.7). This yields a good spectral gap for the diffusion ξ_t restricted to a basin of attraction and mimics the fast convergence of the diffusion to metastable states.
- The second one is a refinement a transport technique mentioned by Chafaï and Malrieu [CM10] leading to a mean-difference estimate (cf. Theorem 1.9). It contains the leading order contribution in ε of the spectral gap of L, resulting from exponential long waiting times of jumps between metastable states of the diffusion.

From these two ingredients, the Eyring-Kramers formula for the first eigenvalue directly follows as a simple consequence (cf. Corollary 1.10) using again the refinement of the two-scale approach.

The remaining part of the introduction is organized as follows: In Section 1.1, we state the precise assumptions on the Hamiltonian H. Whereas in Section 1.2, we state the main results of the article and outline the main idea of the new proof of the Eyring-Kramers formula.

1.1. Assumptions on the Hamiltonian H

Before starting the precise assumptions on the Hamiltonian H, we are introducing the notion of a *Morse function.*

Definition 1.1. A smooth function $H : \mathbb{R}^n \to \mathbb{R}$ is a Morse function, if the Hessian $\nabla^2 H$ of H is non-degenerated on the set of critical points. More precisely, for some $1 \leq C < \infty$ it holds

$$\forall x \in \mathcal{S} := \{x \in \mathbb{R}^n : \nabla H = 0\} : \quad \frac{1}{C} \leq \|\nabla^2 H(x)\| \leq C. \tag{1.5}$$

We make the following assumption on the Hamiltonian H.

Assumption 1.2. *We assume that $H \in C^3(\mathbb{R}^n, \mathbb{R})$ is a Morse function. Further, for some constants $C_H > 0$ and $K_H \geq 0$ it holds*

$$\liminf_{|x|\to\infty} |\nabla H| \geq C_H. \tag{A1}$$

$$\liminf_{|x|\to\infty} |\nabla H|^2 - \Delta H \geq -K_H. \tag{A2}$$

The Assumption 1.2 yields the following consequences for the Hamiltonian H:

- The condition (**A1**) ensures that e^{-H} is integrable and therefore can be normalized to a probability measure on $\mathbb{R}^N$. Hence, the Gibbs measure μ given by (1.4) is well defined.
- A combination of the condition (**A1**) and (**A2**) ensures that there exists a spectral gap for the operator L given by (1.1). For the definition of the spectral gap and for the argument, we refer the reader to the Appendix C. Equivalently, this means by the variational characterization of the spectral gap of L (cf. Lemma 1.5) that the Gibbs measure μ given by (1.4) satisfies a Poincaré inequality for sufficiently small ε.
- Basically, the *Lyapunov-type* condition (**A2**) allows to recover the spectral gap of the full Gibbs measure μ from the spectral gap of the Gibbs measure μ_R restricted to some ball B_R with large but fixed radius $R > 0$ (cf. Section 4.1 or [BCG08]).
- The Morse assumption (1.5) together with the growth condition (**A1**) ensures that the set $\mathcal{S}$ of critical points is discrete and finite. In particular, it follows that the set of local minima $\mathcal{M} = \{m_1, \ldots, m_M\}$ is also finite i.e. $M := \#\mathcal{M} < \infty$.

For the proof of the Eyring-Kramers formula we need an additional non-degeneracy assumption concerning the values of $H(x)$ at critical points $x \in \mathcal{S}$. For that purpose, let us introduce the saddle height $\widehat{H}(m_i, m_j)$ between two local minima m_i, m_j by

$$\widehat{H}(m_i, m_j) = \inf \left\{ \max_{s\in[0,1]} H(\gamma(s)) : \gamma \in C([0,1], \mathbb{R}^n), \gamma(0) = m_i, \gamma(1) = m_j \right\}.$$

Assumption 1.3. *There exists $\delta > 0$ such that:*

(i) The saddle height between two local minima m_i, m_j is attained at a unique critical point $s_{i,j} \in \mathcal{S}$, i.e. it holds $H(s_{i,j}) = \widehat{H}(m_i, m_j)$.
It follows from Assumption 1.2 that $s_{i,j}$ is a saddle point of index one, i.e. $\left\{x \in \mathbb{R}^n : \left\langle \nabla^2 H(s_{i,j})x, x \right\rangle \leq 0\right\}$ is one-dimensional.

(ii) *The set of local minima $\mathcal{M} = \{m_1, \ldots, m_M\}$ is ordered such that m_1 is the global minima and for all $i \in \{3, \ldots, M\}$ yields*

$$H(s_{1,2}) - H(m_2) \geq H(s_{1,i}) - H(m_i) - \delta.$$

The Assumption 1.2 and 1.3 are almost the same as the ones used by [BGK05]. However, there are two minor differences which are outlined in the next remark.

Remark 1.4. We compare our conditions on H with the conditions of [BGK05].

- For convenience, we assume that the domain of H is $\mathbb{R}^n$. As in [BGK05], our argument would also work for any open and connected subset domain $D \subset \mathbb{R}^n$ satisfying $H(x_i) \to \infty$ whenever $x_i \to x \in \partial D$.
- Note that the non-degeneracy assumption (1.5) holds for all critical points x of H. In [BGK05], the assumption (1.5) is only needed for the local minima m_i and the saddles $s_{i,j}$. We need this slightly stronger assumption in the proof of the local Poincaré inequality (cf. Theorem 1.7). In the argument of [BGK05] does not need a local Poincaré inequality.

1.2. Idea of the new proof of the Eyring-Kramers formula and main results

Due to the fact that our approach is motivated by functional inequalities, we are using the well-known variational characterization of the spectral gap by the *Poincaré inequality.*

Lemma 1.5 (Variational characterization of the spectral gap of L)**.** *We say that the Gibbs measure μ on $\mathbb{R}^n$ given by* (1.4) *satisfies the* Poincaré inequality $\mathrm{PI}(\varrho)$ *with constant $\varrho > 0$ if for all functions $f : \mathbb{R}^n \to \mathbb{R}$*

$$\operatorname{var}_\mu(f) := \int \left(f - \int f \mathrm{d}\mu \right)^2 \mathrm{d}\mu \leq \frac{1}{\varrho} \int |\nabla f|^2 \,\mathrm{d}\mu \;\; = \frac{1}{\varrho\varepsilon} \mathcal{E}(f), \qquad \mathrm{PI}(\varrho)$$

where the Dirichlet form $\mathcal{E}$ is given by (1.2)*.*
Then, the operator L given by (1.1) has a spectral gap of size $\varrho\varepsilon$ if and only if the Gibbs measure μ satisfies $\mathrm{PI}(\varrho)$ *with optimal constant ϱ.*

The new proof is based on a refinement of the two-scale approach [GOVW09] and a transport technique introduced in [CM10]. Motivated by the fast convergence of the diffusion ξ_t given by (1.3) to metastable states, we decompose the Gibbs measure μ into a mixture of local Gibbs measures μ_i in the following way. To every local minima $m_i \in \mathcal{M}$ we associate its *basin of attraction* Ω_i defined by

$$\Omega_i := \left\{ y \in \mathbb{R}^n : \lim_{t \to \infty} y_t = m_i, \dot{y}_t = -\nabla H(y_t), y_0 = y \right\}. \tag{1.6}$$

Up to sets of Lebesgue measure zero, the set $\mathcal{P}_M = \{\Omega_i\}_{i=1}^M$ is a partition of $\mathbb{R}^n$. To each element of the partition Ω_i we associate the local Gibbs measure μ_i given by

$$\mu_i(\mathrm{d}x) = \frac{1}{Z_i Z_\mu} \mathbb{1}_{\Omega_i}(x) \; \exp\left(-\frac{H(x)}{\varepsilon}\right) \mathrm{d}x, \qquad \text{where} \quad Z_i = \mu(\Omega_i). \tag{1.7}$$

Note that $\sum_i Z_i = 1$, since $\{\Omega_i\}_{i=1}^M$ is a partition of $\mathbb{R}^n$ and μ a probability measure. The starting point of our argument is a representation of the Gibbs measure μ as a mixture of the mutual singular measures μ_i, namely

$$\mu = Z_1\mu_1 + \cdots + Z_M\mu_M.$$

This decomposition of μ gives rise to a decomposition of the variance $\operatorname{var}_\mu(f)$ into *local variances* and a variance between the averages on Ω_i, which we will call *mean-difference.*

Lemma 1.6. *For all $f : \mathbb{R}^n \to \mathbb{R}$ holds the splitting*

$$\operatorname{var}_\mu(f) = \sum_{i=1}^M Z_i \operatorname{var}_{\mu_i}(f) + \sum_{i=1}^M \sum_{j<i} Z_i Z_j \left(\mathbb{E}_{\mu_i} f - \mathbb{E}_{\mu_j} f\right)^2 . \tag{1.8}$$

We call the term $\operatorname{var}_{\mu_i}(f)$ local variance and $\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2$ mean-difference.

The statement of Lemma (1.6) is verified by an easy straightforward calculation, which we outline for the convenience of the reader in Section 2.

The main results of this paper are good estimates of the right-hand side of (1.8) (cf. Theorem 1.7 and Theorem 1.9 below). The Eyring-Kramer formula is then a simple consequence of these estimates (cf. Corollary 1.10 below).

Let us now turn to the estimation of the local variances $\operatorname{var}_{\mu_i}(f)$. From the heuristic understanding of the process ξ_t given by (1.3), we expect a good behavior of the local Poincaré constant for the local Gibbs measures μ_i, because it would resemble the fast convergence of ξ_t to a neighborhood of the next local minimum. Therefore the local variances should not contribute to the leading order expansion of the total Poincaré constant of μ. This idea is quantified in the following theorem.

Theorem 1.7 (Local Poincaré inequality)**.** *The local measures $\{\mu_i\}_{i=1}^M$, obtained by restricting μ to the basin of attraction Ω_i of the local minima m_i (cf. (1.7)), satisfy* $\mathrm{PI}(\varrho_i)$ *with*

$$\varrho_i^{-1} = O(\varepsilon).$$

Remark 1.8. Using the variational characterization of the spectral gap (cf. Lemma 1.5) one can easily see the following consequence of Theorem 1.7: The spectral gap the diffusion ξ_t given by (1.3) reflected at the boundary of a basin of attraction Ω_i is at least of order 1. This reflects the heuristic idea of a scale separation of the diffusion ξ_t into a fast and a slow scale.

Even if there are simple heuristics for the validity of Theorem 1.7, we need an elaborated machinery for the proof. The reason is that our situation goes beyond the scope of the standard tools for Poincaré inequalities. We outline the argument for Theorem 1.7 in Section 4. It is based on an explicit construction of a perturbation $\tilde{H}$ of the Hamiltonian H (cf. Appendix A).

Let us now turn to the estimation of the mean-difference $\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2$. Note that the term

$$\sum_{i=1}^{M} \sum_{j<i} Z_i Z_j \left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2$$

can be interpreted as the variance of a Bernoulli variable with probabilities $Z_1, \ldots, Z_M$. Therefore we expect to see in the estimation of $\left(\mathbb{E}_{\mu_i} f - \mathbb{E}_{\mu_j} f\right)^2$ the exponential long waiting times of the jumps of the diffusion ξ_t between the basins of attraction Ω_i. We have to estimate the mean-differences on the right hand side of (1.8) in terms of the Dirichlet form $\mathcal{E}$. Practically, we have to find a good upper bound for the constant C in the inequality

$$\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2 \leq C \int |\nabla f|^2 \,\mathrm{d}\mu.$$

Therefore, we introduce in Section 3.1 a new *transport distance* between probability measures, which allows variational characterization of the constant C. By an approximation argument (cf. Section 3.2) we give an explicit construction of a transport interpolation (cf. Section 3.3) estimating asymptotically sharp the constant C, namely

Theorem 1.9 (Mean-difference estimate)**.** *The mean-differences between the measures μ_i and μ_j satisfy*

$$\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2 \leq \frac{Z_\mu}{(2\pi\varepsilon)^{\frac{n}{2}}} \frac{2\pi\varepsilon}{|\lambda^-(s_{i,j})|} \sqrt{|\det \nabla^2 H(s_{i,j})|}\, e^{\frac{1}{\varepsilon} H(s_{i,j})} R(\varepsilon) \int |\nabla f|^2 \,\mathrm{d}\mu,$$

where $\lambda^-(s_{i,j})$ denotes the negative eigenvalue of the Hessian $\nabla^2 H(s_{i,j})$ at the 1-saddle $s_{i,j}$. The multiplicative error term $R(\varepsilon)$ satisfies

$$R(\varepsilon) = 1 + O(\sqrt{\varepsilon}\, |\ln \varepsilon|^{\frac{3}{2}}).$$

The proof of Theorem 1.9 is carried out in full detail in Section 3.

Now, let us turn to the Eyring-Kramers formula. In Section 2 we show that a combination of Theorem 1.7 and Theorem 1.9 with Lemma 1.6 immediately leads to the multidimensional Eyring-Kramers formula (cf. [BGK05, Theorem 1.2]).

Corollary 1.10 (Eyring-Kramers formula)**.** *The measure μ satisfies* PI(ϱ) *with*

$$\frac{1}{\varrho} \leq \frac{2\pi}{|\lambda^-(s_{1,2})|} \frac{\sqrt{|\det \nabla^2 (H(s_{1,2}))|}}{\sqrt{\det \nabla^2 H(m_2)}} e^{\frac{1}{\varepsilon}(H(s_{1,2}) - H(m_2))} \left(1 + O(\sqrt{\varepsilon}\, |\ln \varepsilon|^{\frac{3}{2}})\right). \tag{1.9}$$

In [BGK05, Theorem 1.2]) it is also shown that the upper bound of (1.9) is optimal. In the context of the Arrhenius' law, our new argument for the Eyring-Kramers formula shows some affinity to the proceeding in article [GOV87].

Contents

1. Introduction and main results 1

2. The Eyring-Kramers formula: proof of Corollary 1.10 8

3. The mean-difference estimate: proof of Theorem 1.9 10

4. The local Poincaré inequality: proof of Theorem 1.7 29

A. Construction of the perturbed Hamiltonian $\tilde{H}$: proof of Proposition 4.1 51

B. Gaussian integrals and linear algebra 64

C. Existence of a spectral gap for L 67

D. Conventions and notation 69

2. The Eyring-Kramers formula: proof of Corollary 1.10

The aim of this section is the proof of Corollary 1.10. However, let us first deduce the decomposition result of the variance:

Proof of Lemma 1.6. With the definition of the local Gibbs measure μ_i (cf. (1.7)), we get by straightforward calculation

$$\begin{aligned}\operatorname{var}_\mu(f) &= \int f^2 \mathrm{d}\mu - \left(\int f \mathrm{d}\mu\right)^2 = \sum_i Z_i \int f^2 \mathrm{d}\mu_i - \left(\int f \mathrm{d}\mu\right)^2 \\ &= \sum_i Z_i \underbrace{\left(\int f^2 \mathrm{d}\mu_i - \left(\int f \mathrm{d}\mu_i\right)^2\right)}_{=\operatorname{var}_{\mu_i}(f)} + \left(\sum_i Z_i \left(\int f \mathrm{d}\mu_i\right)^2 - \left(\sum_j Z_j \int f \mathrm{d}\mu_j\right)^2\right).\end{aligned}$$

It only remains to show the identity.

$$\sum_i Z_i \left(\int f \mathrm{d}\mu_i\right)^2 - \left(\sum_j Z_j \int f \mathrm{d}\mu_j\right)^2 = \sum_i \sum_{j<i} Z_i Z_j \left(\int f \mathrm{d}\mu_i - \int f \mathrm{d}\mu_j\right)^2. \tag{2.1}$$

For convenience, let us introduce the notation

$$\bar{f}_i = \int f \mathrm{d}\mu_i.$$

By expanding the square, we get that the left hand side of (2.1) can be written as

$$\sum_i Z_i \bar{f}_i^{\,2} - \sum_{i,j} Z_i Z_j \bar{f}_i \bar{f}_j = \sum_i Z_i (1 - Z_i) \bar{f}_i^{\,2} - 2 \sum_i \sum_{j<i} Z_i Z_j \bar{f}_i \bar{f}_j .$$

By using the identity

$$1 - Z_i = \sum_{j \neq i} Z_j ,$$

the first term on the right hand side of (2.1) can be rewritten as

$$\sum_i Z_i (1 - Z_i) \bar{f}_i^{\,2} = \sum_i \sum_{j<i} Z_i Z_j \bar{f}_i^{\,2} + \sum_i \sum_{j>i} Z_i Z_j \bar{f}_i^{\,2} = \sum_i \sum_{j<i} Z_i Z_j \left(\bar{f}_i^{\,2} + \bar{f}_j^{\,2} \right) .$$

Hence, we finally get the desired identity

$$\begin{aligned} \sum_i Z_i \bar{f}_i^{\,2} - \left(\sum_j Z_j \bar{f}_j \right)^2 &= \sum_i \sum_{j<i} Z_i Z_j (\bar{f}_i^{\,2} - 2 \bar{f}_i \bar{f}_j + \bar{f}_j^{\,2}) \\ &= \sum_i \sum_{j<i} Z_i Z_j \left(\bar{f}_i - \bar{f}_j \right)^2 . \end{aligned}$$

□

Proof of Corollary 1.10. We start from the decomposition of the variance into local variances and mean-differences given by Lemma 1.6. Then an application of Theorem 1.7 and Theorem 1.9 yields the estimate

$$\begin{aligned} \frac{\operatorname{var}_\mu(f)}{\int |\nabla f|^2 \, \mathrm{d}\mu} &\leq \sum_i Z_i \operatorname{var}_{\mu_i}(f) + \sum_i \sum_{j<i} Z_i Z_j \left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f) \right)^2 \\ &\leq O(\varepsilon) + \sum_i \sum_{j<i} \frac{Z_i Z_j Z_\mu}{(2\pi\varepsilon)^{\frac{n}{2}}} \frac{2\pi}{\left|\lambda_H^-(s_{i,j})\right|} \sqrt{|\det \nabla^2 H(s_{i,j})|} e^{\frac{1}{\varepsilon} H(s_{i,j})} R(\varepsilon). \end{aligned} \tag{2.2}$$

There is only left to estimate the product $Z_i Z_j Z_\mu$. An asymptotic expansion of the partition sums Z_μ and Z_{μ_i} shows

$$\begin{aligned} Z_i Z_\mu &= \int_{\Omega_i} e^{-\frac{H}{\varepsilon}} \, \mathrm{d}x = \left(\frac{(2\pi\varepsilon)^{\frac{n}{2}}}{\sqrt{\det \nabla^2 H(m_i)}} + O(\varepsilon^{\frac{n+1}{2}}) \right) e^{-\frac{H(m_i)}{\varepsilon}} , \\ Z_\mu &= \frac{(2\pi\varepsilon)^{\frac{n}{2}}}{\sqrt{\det \nabla^2 H(m_1)}} + O(\varepsilon^{\frac{n+1}{2}}), \end{aligned}$$

where we used the normalization convention $H(m_1) = 0$. Therewith, it holds the asymptotic expansion

$$\frac{Z_i Z_\mu Z_j Z_\mu}{Z_\mu} = \frac{(2\pi\varepsilon)^{\frac{n}{2}} \sqrt{\det \nabla^2 H(m_1)}}{\sqrt{\det \nabla^2 H(m_i) \det \nabla^2 H(m_j)}} e^{-\frac{1}{\varepsilon}(H(m_i) + H(m_j) - H(m_1))} + O(\sqrt{\varepsilon}).$$

Combining this quantity with the mean-difference estimate of Theorem 1.9 leads to

$$\frac{\operatorname{var}_\mu(f)}{\int |\nabla f|^2 \mathrm{d}\mu} \leq \sum_i \sum_{j<i} \frac{2\pi \sqrt{\det \nabla^2 H(m_1) \, |\det \nabla^2 H(s_{i,j})|}}{|\lambda_H^-(s_{i,j})| \sqrt{\det \nabla^2 H(m_i) \det \nabla^2 H(m_j)}} e^{\frac{1}{\varepsilon}(H(s_{i,j})-H(m_i)-H(m_j))} R(\varepsilon) \tag{2.3}$$

Using the property (ii) of Assumption 1.3, we see that the dominating term is given by $i=2$ and $j=1$. This leads to the Kramers-Eyring formula (1.9). ☐

Remark 2.1 (Higher exponential small eigenvalues)*.* The statement of [BGK05, Theorem 1.2]) does not only characterize the second eigenvalue of L (i.e. the spectral gap) but also the higher exponentially small eigenvalues. These characterizations also pop up in our approach: The dominating exponential modes in (2.3) by setting $j=1$ correspond for $i=2,\ldots,M$ to the inverse eigenvalues of L. Hence we believe that our approach should allow to identify these exponentially small eigenvalues. By the variational characterization of eigenvalues of the operator L (cf. Lemma 1.5) one would have to restrict the class of test functions f to the orthogonal complement of the eigenspaces of smaller eigenvalues.

3. The mean-difference estimate: proof of Theorem 1.9

This section is devoted to the proof of Theorem 1.9. We want to estimate the mean-difference $\left(\mathbb{E}_{\mu_i} f - \mathbb{E}_{\mu_j} f\right)^2$ for i and j fixed. The proof consists of four steps:

In the first step, we introduce the *weighted transport distance* in Section 3.1. This distance depends on the transport speed similarly to the Wasserstein distance, but in addition weights the speed of a transported particle w.r.t. the reference measure μ. The weighted transport distance allows in general for a variational characterization of the constant C in the inequality

$$\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2 \leq C \int |\nabla f|^2 \,\mathrm{d}\mu.$$

The problem of finding good estimates of the constant C is then reduced to the problem of finding a good transport between the measures μ_i and μ_j w.r.t. to the weighted transport distance.

For measures as general as μ_i and μ_j, the construction of an explicit transport interpolation is not feasible. Therefore, the second step consists of an approximation, which is done in Section 3.2. There, the restricted measures μ_i and μ_j are replaced by *simpler* measures ν_i and ν_j, namely truncated Gaussians. We show in Lemma 3.6 that this approximation only leads to higher order error terms.

The most import step, the third one, consists of the estimation of the mean-difference w.r.t. the approximations ν_i and ν_j. Because the structure of ν_i and ν_j is very simple, we can explicitly construct a transport interpolation between ν_i and ν_j (see Lemma 3.11 in Section 3.3).

The last step consists of combining the results of Lemma 3.6 and Lemma 3.11 (cf. Section 3.4).

3.1. Mean-difference estimates by transport

At the moment let us consider two arbitrary measures $\nu_0 \ll \mu$ and $\nu_1 \ll \mu$. The overall aim is to estimate the mean-difference

$$\left(\mathbb{E}_{\nu_0}(f) - \mathbb{E}_{\nu_1}(f)\right)^2 .$$

The starting point of the estimation is a representation of the mean-difference as a transport interpolation. This idea goes back to Chafaï and Malrieu [CM10]. However, they used a similar but non-optimal estimate for our purpose. Hence, let us consider a transport map U between ν_0 and ν_1, i.e. the push forward of ν_0 under the map U is given by $U_\sharp \nu_0 = \nu_1$. Further, let $(\Phi_s)_{s\in[0,1]}$ be a smooth interpolation between the identity and the transport map U, i.e.

$$\Phi_0 = \mathrm{Id}, \quad \Phi_1 = U, \quad \text{and} \quad (\Phi_s)_\sharp \nu_0 = \nu_s.$$

The representation of the mean-difference as a transport interpolation is attained by using the fundamental theorem of calculus, i.e.

$$\left(\mathbb{E}_{\nu_0}(f) - \mathbb{E}_{\nu_1}(f)\right)^2 = \left(\int \int_0^1 \frac{\mathrm{d} f(\Phi_s)}{\mathrm{d}s} \, \mathrm{d}s \, \mathrm{d}\nu_0\right)^2 = \left(\int_0^1 \int \langle \nabla f(\Phi_s), \dot{\Phi}_s \rangle \, \mathrm{d}\nu_0 \, \mathrm{d}s\right)^2 .$$

At this point it is tempting to apply the Cauchy-Schwarz inequality in $L^2(\mathrm{d}\nu_0 \times \mathrm{d}s)$ leading to the estimate of Chafaï and Malrieu [CM10]. However, this strategy would not yield the pre-exponential factors in the Eyring-Kramer formula (1.9) (cf. Remark 3.2). On Stephan Luckhaus' advice the authors realized the fact that it really matters on which integral you apply the Cauchy-Schwarz inequality. This insight lead to the following proceeding

$$\begin{aligned}
\left(\mathbb{E}_{\nu_0}(f) - \mathbb{E}_{\nu_1}(f)\right)^2 &= \left(\int_0^1 \int \langle \nabla f(\Phi_s), \dot{\Phi}_s \rangle \, \mathrm{d}\nu_0 \, \mathrm{d}s\right)^2 \\
&= \left(\int_0^1 \int \langle \nabla f, \dot{\Phi}_s \circ \Phi_s^{-1} \rangle \, \mathrm{d}\nu_s \, \mathrm{d}s\right)^2 \\
&\leq \left(\int |\nabla f| \int_0^1 |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \, \mathrm{d}s \, \mathrm{d}\mu\right)^2 \\
&\leq \int \left(\int_0^1 |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \, \mathrm{d}s\right)^2 \mathrm{d}\mu \int |\nabla f|^2 \, \mathrm{d}\mu. \qquad (3.1)
\end{aligned}$$

Note that in the last step we have applied the Cauchy-Schwarz inequality only in $L^2(\mathrm{d}\mu)$ and that the desired Dirichlet integral $\int |\nabla f|^2 \, \mathrm{d}\mu$ is already recovered.

The prefactor in front of the the Dirichlet energy on the right hand side of (3.1) only depends on the transport interpolation $(\Phi_s)_{s\in[0,1]}$. Hence, we can infimize over all possible admissible transport interpolations and arrive at the following definition.

Definition 3.1 (Weighted transport distance)**.** Let μ be an absolutely continuous probability measure on $\mathbb{R}^n$ with connected support. Additionally, let ν_0 and ν_1 be two probability measures such that $\nu_0 \ll \mu$ and $\nu_1 \ll \mu$, then define the *weighted transport distance* by

$$\mathcal{T}_\mu^2(\nu_0,\nu_1) = \inf_{\Phi_s} \int \left(\int_0^1 |\dot\Phi_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \, \mathrm{d}s \right)^2 \mathrm{d}\mu. \tag{3.2}$$

The family $(\Phi_s)_{s\in[0,1]}$ is chosen absolutely continuous in the parameter s such that $\Phi_0 = \mathrm{Id}$ on $\operatorname{supp}\nu_0$ and $(\Phi_1)_\sharp\nu_0 = \nu_1$. For a fixed family and $(\Phi_s)_{s\in[0,1]}$ and a point $x \in \operatorname{supp}\mu$ the *cost density* is defined by

$$\mathcal{A}(x) := \int_0^1 |\dot\Phi_s \circ \Phi_s^{-1}(x)| \, \nu_s(x) \, \mathrm{d}s. \tag{3.3}$$

Remark 3.2 (Relation of $\mathcal{T}_\mu$ to [CM10]). In general, the transport cost $\mathcal{T}_\mu(\nu_0,\nu_1)$ is always smaller than the constant obtained by [CM10, Section 4.6]. Indeed, applying the Cauchy-Schwarz inequality on $L^2(\mathrm{d}s)$ in (3.2) yields

$$\begin{aligned}\mathcal{T}_\mu^2(\nu_0,\nu_1) &\le \inf_{\Phi_s} \int \int_0^1 |\dot\Phi_s \circ \Phi_s^{-1}|^2 \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \, \mathrm{d}s \int_0^1 \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \, \mathrm{d}s \, \mathrm{d}\mu \\ &\le \inf_{\Phi_s} \left(\sup_x \left(\int_0^1 \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu}(x) \, \mathrm{d}s \right) \int \int_0^1 |\dot\Phi_s|^2 \, \mathrm{d}s \, \mathrm{d}\nu_0 \right),\end{aligned}$$

where we used the assumption that $\nu_s \ll \mu$ for all $s \in [0,1]$ in the last $L^1 - L^\infty$ estimate.

Remark 3.3 (Relation of $\mathcal{T}_\mu$ to the L^2-Wasserstein distance W_2). If the support of μ is convex, we can set the transport interpolation $(\Phi_s)_{s\in[0,1]}$ to the linear interpolation map $\Phi_s(x) = (1-s)x + sU(x)$. Assuming that U is the optimal W_2-transport map between ν_0 and ν_1, the estimate in Remark 3.2 becomes

$$\mathcal{T}_\mu^2(\nu_0,\nu_1) \le \left(\sup_x \int_0^1 \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu}(x) \, \mathrm{d}s \right) W_2^2(\nu_0,\nu_1). \tag{3.4}$$

Remark 3.4 (Invariance under time rescaling). The cost density $\mathcal{A}$ given by (3.3) is independent of rescaling the transport interpolation in the parameter s. Indeed, we observe that

$$\mathcal{A}(x) = \int_0^1 |\dot\Phi_s \circ \Phi_s^{-1}(x)| \, \nu_s(x) \, \mathrm{d}s = \int_0^T |\dot\Phi_t^T \circ (\Phi_t^T)^{-1}(x)| \, \nu_t^T(x) \, \mathrm{d}t,$$

where $\Phi_t^T = \Phi_{t/T}$ and $\nu_t^T = \nu_{t/T}$.

In this paragraph, we show that $\mathcal{T}_\mu(\cdot,\cdot)$ actually is a distance justifying the term *weighted transport distance*. It turns out that the distance $\mathcal{T}_\mu(\cdot,\cdot)$ is a metric on a subspace of the space of probability measures on $\mathbb{R}^n$ with finite second moment. The main restriction is that the weighted transport distance is very sensitive to support constraints, which are difficult to check for interpolations between general measures.

Proposition 3.5. *Assume that μ is absolutely continuous w.r.t. the Lebesgue measure. Additionally, assume that μ has convex support and finite second moment. Then $\mathcal{T}_\mu(\cdot,\cdot)$ is a distance on the space*

$$\mathcal{P}_\mu := \{\nu \in \mathcal{P}(\mathbb{R}^n) : \nu \ll \mu,\ \operatorname{supp}\nu \text{ compact}\}.$$

Proof of Proposition 3.5. The symmetry follows from the observation that if $s \mapsto \Phi_s$ is an optimal interpolation between ν_0 and ν_1, then $s \mapsto \Phi_{1-s}$ is an optimal interpolation between ν_1 and ν_0.

Let us consider the definiteness. Therefore, we assume that $\mathcal{T}_\mu(\nu_0,\nu_1) = 0$, then

$$\int_0^1 |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \, \mathrm{d}s = 0, \qquad \mu\text{-a.e.}$$

Hence, by integrating w.r.t. μ and interchanging the order of integration we obtain

$$0 = \int_0^1 \int |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \, \mathrm{d}\mu \, \mathrm{d}s = \int_0^1 \int |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \mathrm{d}\nu_s \, \mathrm{d}s = \int_0^1 \int |\dot{\Phi}_s| \, \mathrm{d}\nu_0 \, \mathrm{d}s.$$

This shows that $\dot{\Phi}_s = 0$ on $\operatorname{supp}\nu_0$. Therefore it holds $\Phi_s = \mathrm{Id}|_{\operatorname{supp}\nu_0}$ resulting in $\nu_s = \nu_0$ for all $s \in [0,1]$.

Let us consider the triangle inequality. We have to show that for arbitrary measures $\nu_0, \nu_{1/2}, \nu_1 \ll \mu$ it holds

$$\mathcal{T}_\mu(\nu_0,\nu_1) \le \mathcal{T}_\mu(\nu_0,\nu_{\frac{1}{2}}) + \mathcal{T}_\mu(\nu_{\frac{1}{2}},\nu_1). \tag{3.5}$$

Let $(\Phi_s)_{s\in[0,1]}$ be an interpolation between ν_0 and ν_1 such that $[0,1] \ni s \mapsto \Phi_{s/2}$ is an interpolation between ν_0 and $\nu_{1/2}$ and $[0,1] \ni s \mapsto \Phi_{s/2+1/2}$ is an interpolation between $\nu_{1/2}$ and ν_1. An application of the triangle inequality in $L^2(\mathrm{d}\mu)$ yields the estimate

$$\begin{aligned}
\mathcal{T}_\mu(\nu_0,\nu_1) &\le \left(\int \left(\int_0^1 |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \mathrm{d}s\right)^2 \mathrm{d}\mu\right)^{\frac{1}{2}} \\
&= \left(\int \left(\int_0^{\frac{1}{2}} |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \mathrm{d}s + \int_{\frac{1}{2}}^1 |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \mathrm{d}s\right)^2 \mathrm{d}\mu\right)^{\frac{1}{2}} \\
&\le \left(\int \left(\int_0^{\frac{1}{2}} |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \mathrm{d}s\right)^2 \mathrm{d}\mu\right)^{\frac{1}{2}} + \left(\int \left(\int_{\frac{1}{2}}^1 |\dot{\Phi}_s \circ \Phi_s^{-1}| \, \frac{\mathrm{d}\nu_s}{\mathrm{d}\mu} \mathrm{d}s\right)^2 \mathrm{d}\mu\right)^{\frac{1}{2}}.
\end{aligned}$$

Because of the invariance of the cost density under rescaling of time (cf. Remark 3.4) and the arbitrariness of the transport $(\Phi_s)_{s\in[0,1]}$, the last inequality already implies the desired triangle inequality (3.5).
In the last step we show that $\mathcal{T}_\mu(\nu_0,\nu_1) < \infty$ for $\nu_0, \nu_1 \in \mathcal{P}_\mu$. For that purpose, we apply the bound in terms of the Wasserstein distance (3.4). Then, it is sufficient to show that $\nu \in \mathcal{P}_\mu$ has finite second moment provided that μ has finite second moment. This fact follows from the change of measure

$$\int |x|^2 \, \mathrm{d}\nu = \int |x|^2 \frac{\mathrm{d}\nu}{\mathrm{d}\mu} \mathrm{d}\mu \le \left\| \frac{\mathrm{d}\nu}{\mathrm{d}\mu} \right\|_{L^\infty} \int |x|^2 \, \mathrm{d}\mu < \infty.$$

In the last inequality we used the observation $\left\| \frac{\mathrm{d}\nu}{\mathrm{d}\mu} \right\|_{L^\infty} < \infty$, which holds due to the compactness of the support of ν. □

3.2. Approximation of the measures μ_i

In this subsection we show that it is sufficient to consider only the mean-difference w.r.t. some auxiliary measures ν_i approximating μ_i. More precisely, the next lemma shows that there are nice measures ν_i which are close to the measures μ_i in the sense of the mean-difference.

Lemma 3.6 (Mean-difference of approximation)**.** *Let ν_i be a truncated Gaussian centered around the local minima m_i with covariance matrix $\Sigma_i = (\nabla^2 H(m_i))^{-1}$, more precisely*

$$\nu_i(\mathrm{d}x) = \frac{1}{Z_{\nu_i}} \, e^{-\frac{\Sigma_i^{-1}[x-m_i]}{2\varepsilon}} \, \mathbb{1}_{E_i}(x) \, \mathrm{d}x, \quad \textit{where} \quad Z_{\nu_i} = \int_{E_i} e^{-\frac{\Sigma_i^{-1}[x-m_i]}{2\varepsilon}} \, \mathrm{d}x. \tag{3.6}$$

Here and further on, we use the convention that for a matrix M and a vector x we write

$$M[x] := \langle x, Mx \rangle \, .$$

The restriction E_i is given by an ellipsoid

$$E_i = \{x \in \mathbb{R}^n : |\Sigma_i^{-\frac{1}{2}}(x - m_i)| \le \sqrt{2\varepsilon}\, \omega(\varepsilon)\}. \tag{3.7}$$

Additionally, assume that μ_i satisfies $\mathrm{PI}(\varrho_i)$ with $\varrho_i^{-1} = O(\varepsilon)$.
Then the following estimate holds

$$(\mathbb{E}_{\nu_i}(f) - \mathbb{E}_{\mu_i}(f))^2 \le O(\varepsilon^{\frac{3}{2}} \omega^3(\varepsilon)) \int |\nabla f|^2 \, \mathrm{d}\mu, \tag{3.8}$$

where the function $\omega(\varepsilon) : \mathbb{R}^+ \to \mathbb{R}^+$ in (3.7) and (3.8) is smooth and monotone satisfying

$$\omega(\varepsilon) \ge |\ln \varepsilon|^{\frac{1}{2}} \qquad \textit{for } \varepsilon < 1.$$

The first step towards the proof of Lemma 3.6 is the following Lemma.

Lemma 3.7. *Consider an arbitrary measure ν_i satisfying $\nu_i \ll \mu_i$. Additionally, assume that μ_i satisfies $\mathrm{PI}(\varrho_i)$ for some $\varrho_i > 0$. Then the following estimate holds*

$$\left(\mathbb{E}_{\nu_i}(f) - \mathbb{E}_{\mu_i}(f)\right)^2 \le \frac{1}{\varrho_i} \operatorname{var}_{\mu_i}\left(\frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\right) \int |\nabla f|^2 \,\mathrm{d}\mu_i. \tag{3.9}$$

Proof of Lemma 3.7. The result is a consequence from the representation of the mean-difference as a covariance. Therefore, we note that $\mathrm{d}\nu_i = \frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\,\mathrm{d}\mu_i$ since $\nu_i \ll \mu_i$ and use the Cauchy-Schwarz inequality for the covariance

$$\begin{aligned}\left(\mathbb{E}_{\nu_i}(f) - \mathbb{E}_{\mu_i}(f)\right)^2 &= \int f \mathrm{d}\nu_i - \int f \mathrm{d}\mu_i = \int f\, \frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\,\mathrm{d}\mu_i - \int f \mathrm{d}\mu_i \underbrace{\int \frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\,\mathrm{d}\mu_i}_{=1} \\ &= \operatorname{cov}^2_{\mu_i}\left(\frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}, f\right) \le \operatorname{var}_{\mu_i}\left(\frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\right) \operatorname{var}_{\mu_i}(f).\end{aligned}$$

Using the fact that μ_i satisfies a Poincaré inequality results in (3.9). ☐

The above lemma tells us that we only need to construct ν_i, which approximates μ_i in variance. The following lemma provides exactly this.

Lemma 3.8 (Approximation in variance)**.** *Assume that the measures ν_i are given by Lemma 3.6. Then the partition sum Z_{ν_i} satisfies for ε small enough*

$$Z_{\nu_i} = (2\pi\varepsilon)^{\frac{n}{2}} \sqrt{\det \Sigma_i}\, \left(1 + O(\sqrt{\varepsilon})\right). \tag{3.10}$$

Additionally, ν_i is a good approximation in variance of μ_i, i.e.

$$\operatorname{var}_{\mu_i}\left(\frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\right) = O(\sqrt{\varepsilon}\, \omega^3(\varepsilon)). \tag{3.11}$$

Proof of Lemma 3.8. The proof of (3.10) reduces to an estimate of a Gaussian integral on the complementary domain $\mathbb{R}^n \backslash E_i$. By recalling, that $(2\pi\varepsilon)^{\frac{n}{2}} \sqrt{\det \Sigma_i}$ is the normalization for a Gaussian with covariance matrix Σ_i, we arrive at

$$Z_{\nu_i} = \int_{E_i} e^{-\frac{\Sigma_i^{-1}[x-m_i]}{2\varepsilon}} \mathrm{d}x = (2\pi\varepsilon)^{\frac{n}{2}} \sqrt{\det \Sigma_i} \left(1 - \frac{1}{(2\pi\varepsilon)^{\frac{n}{2}} \sqrt{\det \Sigma_i}} \int_{\mathbb{R}^n \backslash E_i} e^{-\frac{\Sigma_i^{-1}[x-m_i]}{2\varepsilon}} \mathrm{d}x\right).$$

The integral on the complementary domain $\mathbb{R}^n \backslash E_i$ evaluates by the change of variables $x \mapsto y = (2\varepsilon\Sigma_i)^{-\frac{1}{2}}(x - m_i)$ to

$$\begin{aligned}\frac{1}{(2\pi\varepsilon)^{\frac{n}{2}} \sqrt{\det \Sigma_i}} \int_{\mathbb{R}^n \backslash E_i} e^{-\frac{\Sigma_i^{-1}[x-m_i]}{2\varepsilon}} \mathrm{d}x &= \frac{1}{\pi^{\frac{n}{2}}} \int_{\mathbb{R}^n \backslash B_{\omega(\varepsilon)}} e^{-y^2} \mathrm{d}y = \frac{n}{\Gamma(\frac{n}{2}+1)} \int_{\omega(\varepsilon)}^{\infty} r^{n-1} e^{-r^2} \mathrm{d}r \\ &= \frac{1}{\Gamma(\frac{n}{2})} \int_{\omega^2(\varepsilon)}^{\infty} r^{\frac{n}{2}-1} e^{-r} \mathrm{d}r = \frac{\Gamma(\frac{n}{2}, \omega^2(\varepsilon))}{\Gamma(\frac{n}{2})},\end{aligned}$$

where $\Gamma(\frac{n}{2}, \omega^2(\varepsilon))$ is the complementary incomplete Gamma function. It has the asymptotic expansion [Olv97, p. 109-112] given by

$$\Gamma\left(\frac{n}{2}, \omega^2(\varepsilon)\right) = O(e^{-\omega^2(\varepsilon)}\omega^{n-2}(\varepsilon)), \quad \text{for } \omega(\varepsilon) \geq \sqrt{n}.$$

We obtain (3.10) by the choice of $\omega(\varepsilon) \geq |\ln \varepsilon|^{\frac{1}{2}}$, since the error becomes

$$O(e^{-\omega^2(\varepsilon)}\omega^{n-2}(\varepsilon)) = O(\varepsilon\, |\ln \varepsilon|^{\frac{n}{2}-1}) = O(\sqrt{\varepsilon}), \qquad \text{for } \varepsilon \leq e^{-n}.$$

For the proof of (3.11), we compare the asymptotic expression for $Z_{\mu_i} = Z_i Z_\mu e^{\varepsilon^{-1} m_i}$ from (2.2) and Z_{ν_i} and obtain

$$Z_{\mu_i} = Z_{\nu_i} + O(\sqrt{\varepsilon}). \tag{3.12}$$

The relative density of ν_i w.r.t. μ_i can be estimates by a Taylor expansion of H around m_i. By the definition of ν_i given in (3.6), we obtain that $\Sigma_i^{-1}[y - m_i] - H_i(y) = O(|y - m_i|^3)$. This observation together with (3.12) leads to

$$\frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}(y) = \frac{Z_{\mu_i}}{Z_{\nu_i}} e^{-\frac{1}{2\varepsilon}\Sigma_i^{-1}[y-m_i] + \frac{1}{2\varepsilon}H_i(y)} \mathbb{1}_{E_i}(y) = \frac{Z_{\mu_i}}{Z_{\nu_i}} e^{\frac{O(|y-m_i|^3)}{\varepsilon}} \mathbb{1}_{E_i}(y) = 1 + O(\sqrt{\varepsilon}\, \omega^3(\varepsilon)).$$

Now, the conclusion directly follows from the definition of the variance

$$\begin{aligned} \operatorname{var}_{\mu_i}\left(\frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\right) &= \int_{E_i} \left(\frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\right)^2 \mathrm{d}\mu_i - \left(\int \frac{\mathrm{d}\nu_i}{\mathrm{d}\mu_i}\, \mathrm{d}\mu_i\right)^2 \\ &= \int_{E_i} 1 + O(\sqrt{\varepsilon}\, \omega^3(\varepsilon))\, \mathrm{d}\mu_i - \left(\int_{E_i} \mathrm{d}\nu_i\right)^2 = 1 + O(\sqrt{\varepsilon}\, \omega^3(\varepsilon)) - 1. \end{aligned}$$

□

Proof of Lemma 3.6. A combination of Lemma 3.7 and Lemma 3.8 together with the assumption $\varrho_i^{-1} = O(\varepsilon)$ immediately reveals

$$(\mathbb{E}_{\nu_i}(f) - \mathbb{E}_{\mu_i}(f))^2 \overset{(3.9),(3.11)}{\leq} O(\varepsilon^{\frac{3}{2}}\omega^3(\varepsilon)) \int |\nabla f|^2\, \mathrm{d}\mu_i.$$

□

3.3. Affine transport interpolation

The aim of this subsection is to estimate $\left(\mathbb{E}_{\nu_i}(f) - \mathbb{E}_{\nu_j}(f)\right)^2$ with the help of the weighted transport distance $\mathcal{T}_\mu(\nu_i, \nu_j)$ introduced in Section 3.1. The main result of this section estimates the weighted transport distance $\mathcal{T}_\mu(\nu_i, \nu_j)$ and is formulated in Lemma 3.11. For the proof of Lemma 3.11, we construct an explicit transport interpolation between

ν_i and ν_j w.r.t. the measure μ. We start with a class of possible transport interpolations and optimize the weighted transport cost in this class.

Now, we state the main idea of this optimization procedure. Recall that the measures ν_i and ν_j are truncated Gaussians by the approximation we have done in the previous Section 3.2. Hence, the measures ν_i and ν_j are characterized by their mean and covariance matrix. We will choose the transport interpolation (cf. Section 3.3.1) such that the push forward measures $\nu_s := (\Phi_s)_\sharp \nu_0$ are again truncated Gaussians. Hence, it is sufficient to optimize among all paths γ connecting the minima m_i and m_j and all covariance matrices interpolating between Σ_i and Σ_j.

3.3.1. Definition of regular affine transport interpolations

Let us state in this section the class of transport interpolation among we want to optimize the weighted transport cost.

Definition 3.9 (Affine transport interpolations)**.** Assume that the measures ν_i and ν_j are given by Lemma 3.6. More precisely, $\nu_i = \mathcal{N}(m_i, \varepsilon^{-1}\Sigma_i)\llcorner E_i$ and $\nu_j = \mathcal{N}(m_j, \varepsilon^{-1}\Sigma_j)\llcorner E_j$ are truncated Gaussians centered in m_i and m_j with covariance matrices $\varepsilon^{-1}\Sigma_i$ and $\varepsilon^{-1}\Sigma_j$. The restriction E_i and E_j are given for $l = 1, 2$ by the ellipsoid

$$E_l = \{x \in \mathbb{R}^n : |\Sigma_l^{-\frac{1}{2}}(x - m_l)| \leq \sqrt{2\varepsilon}\, \omega(\varepsilon)\}, \qquad \text{where } \omega(\varepsilon) \geq |\ln \varepsilon|^{\frac{1}{2}} .$$

A transport interpolation Φ_s between ν_i and ν_j is called *affine transport interpolation* if there exists

- an interpolation path $(\gamma_s)_{s\in[0,T]}$ between $m_i = \gamma_0$ and $m_j = \gamma_T$ satisfying
$$\gamma = (\gamma_s)_{s\in[0,T]} \in C^2([0,T], \mathbb{R}^n) \qquad \text{and} \qquad \forall s \in [0,T] : \dot\gamma_s \in S^{n-1}, \tag{3.13}$$
- an interpolation path $(\Sigma_s)_{s\in[0,T]}$ of covariance matrices between Σ_i and Σ_j satisfying
$$\Sigma = (\Sigma_s)_{s\in[0,T]} \in C^2([0,T], S^+(n)), \quad \Sigma_0 = \Sigma_i \quad \text{and} \quad \Sigma_T = \Sigma_j,$$

such that the transport interpolation $(\Phi_s)_{s\in[0,T]}$ is given by

$$\Phi_s(x) = \Sigma_s^{\frac{1}{2}} \Sigma_0^{-\frac{1}{2}}(x - m_0) + \gamma_s. \tag{3.14}$$

Since the cost density $\mathcal{A}$ given by (3.3) is invariant under rescaling of time (cf. Remark 3.4), one can always assume that the interpolation path γ_s is parameterized by arc-length. Hence, the condition $\dot\gamma_s \in S^{n-1}$ (cf. (3.13)) is not restricting.

We want to emphasize that for an affine transport interpolation $(\Phi_s)_{s\in[0,T]}$ the push forward measure $(\Phi_s)_\sharp\nu_0 = \nu_s$ is again a truncated Gaussian $\mathcal{N}(\gamma_s, \varepsilon^{-1}\Sigma_s)\llcorner E_s$, where E_s is the support of ν_s being again an ellipsoid in $\mathbb{R}^n$ given by

$$E_s = \{x \in \mathbb{R}^n : |\Sigma_s^{-\frac{1}{2}}(x-\gamma_s)| \le \sqrt{2\varepsilon}\,\omega(\varepsilon)\}. \tag{3.15}$$

Therewith, the partition sum of ν_s is given by (cf. (3.10))

$$Z_{\nu_s} = (2\pi\varepsilon)^{\frac{n}{2}}\sqrt{\det\Sigma_s}(1+O(\sqrt{\varepsilon})). \tag{3.16}$$

By denoting $\sigma_s = \Sigma_s^{\frac{1}{2}}$ and using the definition (3.14) of the affine transport interpolation $(\Phi_s)_{s\in[0,T]}$, we arrive at the relations

$$\begin{aligned} \dot\Phi_s(x) &= \dot\sigma_s\sigma_0^{-1}(x-m_0)+\dot\gamma_s, \\ \Phi_s^{-1}(y) &= \sigma_0\sigma_s^{-1}(y-\gamma_s)+m_0, \\ \dot\Phi_s\circ\Phi_s^{-1}(y) &= \dot\sigma_s\sigma_s^{-1}(y-\gamma_s)+\dot\gamma_s. \end{aligned}$$

Among all possible affine transport interpolations we are considering only those satisfying the following regularity assumption.

Assumption 3.10 (Regular affine transport interpolations)**.** *The affine transport interpolation $(\gamma_s,\Sigma_s)_{s\in[0,T]}$ belongs to the class regular affine transport interpolations if the length $T < T^*$ is bounded by some uniform $T^* > 0$ large enough. Further, for a uniform constant $c_\gamma > 0$ it holds*

$$\inf\{r(x,y,z) : x,y,z\in\gamma, x\neq y\neq z\neq x\} \ge c_\gamma, \tag{3.17}$$

where $r(x,y,z)$ denotes the radius of the unique circle through the three distinct points x, y and z. Furthermore, there exists a uniform constant $C_\Sigma \ge 1$ for which

$$C_\Sigma^{-1}\,\mathrm{Id} \le \Sigma_s \le C_\Sigma\,\mathrm{Id} \qquad \text{and} \qquad \|\dot\Sigma_s\| \le C_\Sigma. \tag{3.18}$$

The infimum in condition (3.17) is called *global radius of curvature* (cf. [GMSvdM02]). It ensures that a small neighborhood of size $\frac{c_\gamma}{2}$ around γ is not self-intersecting, since the infimum can only be attained for the following three cases:

(i) All three points in a minimizing sequence of (3.17) coalesce to a point at which the radius of curvature is minimal.

(ii) Two points coalesce to a single point and the third converges to another point, such that the both points are a pair of closest approach.

(iii) Two points coalesce to a single point and the third converges to the starting or ending point of γ.



In the following calculations, there often occurs a multiplicative error of the form $1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))$. Therefore let us introduce for convenience the notation "$\approx$" meaning "$=$" up to the multiplicative error $1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))$. The symbols "$\lesssim$" and "$\gtrsim$" have the analogous meaning.

Now, we can formulate the key ingredient for the proof of Theorem 1.9, namely the estimation of the weighted transport distance $\mathcal{T}_\mu(\nu_i, \nu_j)$.

Lemma 3.11. *Assume that ν_i and ν_j are given by Lemma 3.6. Then the weighted transport distance $\mathcal{T}_\mu(\nu_i, \nu_j)$ can be estimated as*

$$\begin{aligned}\mathcal{T}_\mu^2(\nu_0, \nu_1) &= \inf_{\Phi_s} \int \left(\int_0^1 |\dot{\Phi}_s \circ \Phi_s^{-1}| \,\frac{\mathrm{d}\nu_s}{\mathrm{d}\mu}\,\mathrm{d}s \right)^2 \mathrm{d}\mu \\ &\leq \inf_{\Psi_s} \int \left(\int_0^1 |\dot{\Psi}_s \circ \Psi_s^{-1}| \,\frac{\mathrm{d}\nu_s}{\mathrm{d}\mu}\,\mathrm{d}s \right)^2 \mathrm{d}\mu \\ &\lesssim \frac{Z_\mu}{(2\pi\varepsilon)^{\frac{n}{2}}}\, e^{\frac{H(s_{i,j})}{\varepsilon}}\, 2\pi\varepsilon \left(\frac{\sqrt{|\det(\nabla^2 H(s_{i,j})|}}{|\lambda^-|} + \frac{T(C_\Sigma)^{\frac{n-1}{2}}}{\sqrt{2\pi\varepsilon}} e^{-\omega^2(\varepsilon)} \right), \end{aligned} \tag{3.19}$$

where the infimum over Ψ_s only considers regular affine transport interpolations Ψ_s in the sense of Assumption 3.10.
In particular, if we choose $\omega(\varepsilon) \geq |\ln\varepsilon|^{\frac{1}{2}}$, which is enforced by Lemma 3.6, we get the estimate

$$\mathcal{T}_\mu^2(\nu_i, \nu_j) \leq \frac{Z_\mu}{(2\pi\varepsilon)^{\frac{n}{2}}} \frac{2\pi\varepsilon}{|\lambda^-|} \sqrt{|\det(\nabla^2 H(s_{i,j})|}\, e^{\frac{H(s_{i,j})}{\varepsilon}} \left(1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))\right). \tag{3.20}$$

Remark 3.12. With a little bit of extra work one could show that the estimate (3.19) in fact is optimal in the class of regular affine transport interpolations Ψ_s in the sense of Assumption 3.10 up to a multiplicative error of order $1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))$. However, we omitted the argument because it would not provide any further insights for the main goal of this article, namely a new proof of the Eyring-Kramer formula (1.9).

Because the proof of Lemma 3.11 represents the core of our new proof of the Eyring-Kramer formula (1.9), we state the argument in full detail. It consits of three steps carried out in the following sections:

- In Section 3.3.2, we carry out some preparatory work: We introduce tube coordinates on the support of the transport cost $\mathcal{A}$ given by (3.3) (cf. Lemma 3.13), we deduce a pointwise estimate on the transport cost $\mathcal{A}$, and we give a rough a priori estimate on the transport cost $\mathcal{A}$.
- In Section 3.3.3, we split the transport cost into a transport cost around the saddle and the complement. We also estimate the transport cost of the complement yielding the second summand in the desired estimate (3.19).
- In Section 3.3.4, we finally deduce a sharp estimate of the transport cost around the saddle yielding the first summand in the desired estimate (3.19).

3.3.2. Preparations for the proof of Lemma 3.11

The main reason for making the regularity Assumption 3.10 on affine transport interpolations is that then one can introduce tube coordinates around the path γ. In these coordinates, the calculation of the cost density $\mathcal{A}$ given by (3.3) becomes a lot handier.

We start with defining the caps E_0^- and E_T^+ as

$$E_0^- := \{x \in E_0 : \langle x - \gamma_0, \dot{\gamma}_0\rangle < 0\} \quad \text{and} \quad E_T^+ := \{x \in E_T : \langle x - \gamma_T, \dot{\gamma}_T\rangle > 0\},$$

The caps E_0^- and E_T^+ have no contribution to the total cost but unfortunately need some special treatment. Further, we define the slices V_s with $s \in [0,T]$

$$V_s = \{x \in \operatorname{span}\{\dot{\gamma}_s\}^\perp : |\Sigma_s^{-\frac{1}{2}} x| \le \sqrt{2\varepsilon}\,\omega(\varepsilon)\}$$

In $\operatorname{span} V_s$ we can choose a basis $e_s^2, \dots, e_s^n$ smoothly depending on the parameter s. Especially there exists a family of rotational matrices $(Q_s)_{s\in[0,T]} \in C^2([0,T], SO(n))$ satisfying the same regularity assumption as the family $(\Sigma_\tau)_{\tau\in[0,T]}$ such that

$$Q_s e^1 = \dot{\gamma}_s, \qquad Q_s e^i = e_s^i, \quad \text{for } i = 2, \dots, n, \tag{3.21}$$

where $(e^1, \dots, e^n)$ is the canonical basis of $\mathbb{R}^n$.

Let use now define the tube E as

$$E = \bigcup_{s\in[0,T]} (\gamma_s + V_s)\,.$$

The support of the cost density $\mathcal{A}$ given by (3.3) is now given by

$$\operatorname{supp}\mathcal{A} = \bigcup_{s\in[0,T]} E_s = E_0^- \cup E \cup E_T^+. \tag{3.22}$$

By the definition (3.15) of E_s and the uniform bound (3.18) on Σ_s it holds

$$\operatorname{diam} V_s \le 2\sqrt{2\varepsilon C_\Sigma}\,\omega(\varepsilon). \tag{3.23}$$

Therewith, we find

$$\operatorname{supp}\mathcal{A} \subset B_{2\sqrt{2\varepsilon C_\Sigma}\,\omega(\varepsilon)}((\gamma_\tau)_{\tau\in[0,T]}) := \left\{x \in \mathbb{R}^n : |x - \gamma_\tau| \le 2\sqrt{2\varepsilon C_\Sigma}\,\omega(\varepsilon)\right\}.$$

The assumption (3.15) ensures that $B_{2\sqrt{2\varepsilon C_\Sigma}\,\omega(\varepsilon)}((\gamma_\tau)_{\tau\in[0,T]})$ is not self-intersecting for any ε small enough. The next lemma just states that by changing to tube coordinates in E one can asymptotically neglect the Jacobian determinant $\det J$.

Lemma 3.13 (Change of coordinates)**.** *For the change of coordinates $(\tau, z) \mapsto x = \gamma_\tau + z_\tau$ with $z_\tau \in V_\tau$ it holds for any function ξ on E that*

$$\int_E \xi(x)\,\mathrm{d}x \approx \int_0^T \int_{V_\tau} \xi(\gamma_\tau + z_\tau)\,\mathrm{d}z_\tau\,\mathrm{d}\tau.$$

Proof of Lemma 3.13. We use the representation of the tube coordinates via (3.21). Therewith it holds that $x = \gamma_\tau + Q_\tau z$, where $z \in \{0\} \times \mathbb{R}^{n-1}$. Then, the Jacobian J of the coordinate change $x \mapsto (\tau, Q_\tau z)$ is given by

$$J = (\dot\gamma_\tau + \dot Q_\tau z, (Q_\tau)_2, \dots, (Q_\tau)_n) \in \mathbb{R}^{n\times n},$$

where $(Q_\tau)_i$ denotes the i-th column of Q_τ. By the definition (3.21) of Q_τ follows $\dot\gamma_\tau = (Q_\tau)_1$. Hence, we have the representation $J = Q_\tau + \dot Q_\tau z \otimes e_1$. The determinant of J is then given by

$$\det J = \det\Big(Q_\tau + \dot Q_\tau z \otimes e_1\Big) = \underbrace{\det(Q_\tau)}_{=1} \det\Big(\mathrm{Id} + (Q_\tau^\top \dot Q_\tau z) \otimes e_1\Big) = 1 + \Big(Q_\tau^\top \dot Q_\tau z\Big)_1 .$$

By Assumption 3.10 it holds $\|\dot Q_\tau\| \le C_\Sigma$, from which we conclude $(Q_\tau^\top \dot Q_\tau z)_{1,1} = O(z)$. Since $Q_\tau z \in V_\tau$, we get $O(z) = O(\sqrt{\varepsilon}\,\omega(\varepsilon))$ by (3.23). Hence we get

$$\det J = 1 + O(\sqrt{\varepsilon}\,\omega(\varepsilon)),$$

which concludes the proof. ☐



Figure 1: The support of $\mathcal{A}$ in tube coordinates.

An important tool is the following auxiliary estimate.

Lemma 3.14 (Pointwise estimate of the cost-density $\mathcal{A}$)**.** *For $x \in \operatorname{supp}\mathcal{A}$ we define*

$$\tau = \operatorname*{arg\,min}_{s\in[0,T]} |x - \gamma_s| \qquad \text{and} \qquad z_\tau := x - \gamma_\tau. \tag{3.24}$$

Then the following estimate holds

$$\mathcal{A}(x) \lesssim (2\pi\varepsilon)^{-\frac{n-1}{2}} \sqrt{\det\nolimits_{1,1}(Q_\tau^\top \tilde\Sigma_\tau^{-1} Q_\tau)}\;\; e^{-\frac{\tilde\Sigma_\tau^{-1}[z_\tau]}{2\varepsilon}} =: P_\tau\; e^{-\frac{\tilde\Sigma_\tau^{-1}[z_\tau]}{2\varepsilon}}, \tag{3.25}$$

where Q_τ is defined in (3.21) and $\tilde\Sigma_\tau^{-1}$ is given by

$$\tilde\Sigma_\tau^{-1} = \Sigma_\tau^{-1} - \frac{1}{\Sigma_\tau^{-1}[\dot\gamma_\tau]} \Sigma_\tau^{-1}\dot\gamma_\tau \otimes \Sigma_\tau^{-1}\dot\gamma_\tau. \tag{3.26}$$

Further, $\det_{1,1} A$ is the determinant of the matrix obtained from A removing the first row and column.

Remark 3.15. With a little bit of additionally work, one could show that (3.25) holds with "$\approx$" instead of "$\lesssim$". It follows from (3.26) that the matrix $\tilde{\Sigma}_\tau^{-1}$ is positive definite. Hence, $\mathcal{A}$ is an $\mathbb{R}^{n-1}$-dimensional Gaussian on the slice $\gamma_\tau + V_\tau$ up to approximation errors.

Proof of Lemma 3.24. We start the proof with some preliminary remarks and results. By the regularity Assumption 3.10 on the transport interpolation, we find that for all $x \in \operatorname{supp} \mathcal{A}$ holds uniformly

$$I_T(x) := \{s : E_s \ni x\} \quad \text{satisfies} \quad \mathcal{H}^1(I_T(x)) = O(\sup_{s\in[0,T]} \operatorname{diam}(E_s)) = O(\sqrt{\varepsilon}\,\omega(\varepsilon)).$$

This allows to linearize the transport interpolation around τ given in (3.24). It holds for s such that $x \in E_s$

$$\begin{aligned} \Sigma_s^{-1}[x-\gamma_s] &= \Sigma_\tau^{-1}[\gamma_\tau + z_\tau - \gamma_s] + O(\varepsilon^{\frac{3}{2}}\omega^3(\varepsilon)) \\ &= \Sigma_\tau^{-1}[(\tau - s)\dot{\gamma}_\tau + z_\tau] + O(\varepsilon^{\frac{3}{2}}\omega^3(\varepsilon)). \end{aligned} \tag{3.27}$$

For similar reasons, we can linearize the determinant $\det \Sigma_s$ and have $\det\Sigma_s = \det\Sigma_\tau + O(\sqrt{\varepsilon}\,\omega(\varepsilon))$. Finally, we have the following bound on the transport speed

$$\begin{aligned} |\dot{\Phi}_s \circ \Phi_s^{-1}(x)| \mathbb{1}_{E_s}(x) &= \left|\dot{\sigma}_s\sigma_s^{-1}(x-\gamma_s) + \dot{\gamma}_s\right| \mathbb{1}_{E_s}(x) \\ &\le \left(\left|\dot{\sigma}_s\sigma_s^{-1}(x-\gamma_s)\right| + |\dot{\gamma}_s|\right) \mathbb{1}_{E_s}(x) \\ &\le (C_\Sigma\,|x-\gamma_s| + 1)\,\mathbb{1}_{E_s}(x) = \left(1 + O(\sqrt{\varepsilon}\,\omega(\varepsilon))\right)\mathbb{1}_{E_s}(x). \end{aligned} \tag{3.28}$$

Let us first consider the case $x \in E$. We use (3.16), (3.27) and (3.28) to arrive with $x = \gamma_\tau + z_\tau$ where $z_\tau \in V_\tau$ at

$$\begin{aligned} \mathcal{A}(x) &= \int_{I_T(x)} |\dot{\Phi}_s \circ \Phi_s^{-1}(x)| \, \frac{1}{Z_{\nu_s}} \exp\left(-\frac{1}{2\varepsilon}\Sigma_s^{-1}[x-\gamma_s]\right) \mathbb{1}_{E_s}(x)\,\mathrm{d}s \\ &\le \frac{1}{(2\pi\varepsilon)^{\frac{n}{2}}} \int_{I_T(x)} \frac{1 + O(\sqrt{\varepsilon}\,\omega(\varepsilon))}{\sqrt{\det\Sigma_s}} \exp\left(-\frac{1}{2\varepsilon}\Sigma_s^{-1}[x-\gamma_s]\right)\mathrm{d}s \\ &\le \frac{1}{(2\pi\varepsilon)^{\frac{n}{2}}\sqrt{\det\Sigma_\tau}} \int_{\mathbb{R}} \exp\left(-\frac{1}{2\varepsilon}\Sigma_\tau^{-1}[(\tau-s)\dot{\gamma}_\tau + z_\tau]\right)\mathrm{d}s\,\left(1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))\right) \\ &= \frac{\sqrt{\det\Sigma_\tau^{-1}}}{(2\pi\varepsilon)^{\frac{n}{2}}} \frac{\sqrt{2\pi\varepsilon}}{\sqrt{\Sigma_\tau^{-1}[\dot{\gamma}_\tau]}} \exp\left(-\frac{1}{2\varepsilon}\tilde{\Sigma}_\tau^{-1}[z_\tau]\right)\left(1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon)\right), \end{aligned}$$

where the last step follows by an application of a partial Gaussian integration (cf. Lemma B.1). Finally, by using the relation (B.2), we get that

$$\frac{\Sigma_\tau^{-1}}{\Sigma_\tau^{-1}[\dot{\gamma}_\tau]} = \det\nolimits_{1,1}(Q_\tau^\top \tilde{\Sigma}_\tau^{-1} Q_\tau),$$

and conclude the hypothesis for this case.

Let us now consider the case $x \in E_0^- \cup E_T^+$. For convenience, we only consider the case $x \in E_0^-$. By the definition of E_0^- it holds $\tau = 0$. The integration domain $I_T(x)$ is now given by

$$I_T(x) = [0, s^*) \quad \text{with} \quad s^* = O(\sqrt{\varepsilon}\,\omega(\varepsilon)). \tag{3.29}$$

Therewith, we can estimate $\mathcal{A}(x)$ in the same way as in the case of $x \in E$ and conclude the proof. ☐

We only need one more ingredient for the proof of Lemma 3.11. It is a a priori estimate on the cost density $\mathcal{A}$.

Lemma 3.16. *The following estimates for $\mathcal{A}$ hold:*

$$\int \mathcal{A}(x)\,\mathrm{d}x \lesssim T, \qquad \textit{and} \tag{3.30}$$

$$\mathcal{A}(x) \lesssim \left(\frac{C_\Sigma}{2\pi\varepsilon}\right)^{\frac{n-1}{2}} \qquad \textit{for } x \in \operatorname{supp}\mathcal{A}. \tag{3.31}$$

Proof of Lemma 3.16. Let us first consider the estimate (3.30). It follows from the characterization (3.22) of the support of $\mathcal{A}$ that

$$\int \mathcal{A}(x)\,\mathrm{d}x = \int_E \mathcal{A}(x)\,\mathrm{d}x + \int_{E_0^- \cup E_T^+} \mathcal{A}(x)\,\mathrm{d}x. \tag{3.32}$$

Now, we estimate the first term on the right hand side of the last identity. Using the change to tube coordinates of Lemma 3.13 and noting that the upper bound (3.25) is a Gaussian density on V_τ for $\tau \in [0, T]$, we can easily infer that

$$\int_E \mathcal{A}(x)\,\mathrm{d}x \lesssim |\gamma| = T$$

Let us turn to the second term on the right hand side of (3.32). For convenience, we only consider the integral w.r.t. the cap E_0^-. It follows from (3.28) and (3.29) that

$$\begin{aligned}\int_{E_0^-} \mathcal{A}(x)\,\mathrm{d}x &\lesssim \int_{E_0^-}\int_0^1 \nu_s(x)\,\mathrm{d}s\,\mathrm{d}x = \int_0^{s^*}\int_{E_0^-} \nu_s(x)\,\mathrm{d}s\,\mathrm{d}x \\ &\lesssim \int_0^{s^*}\int \nu_s(x)\,\mathrm{d}s\,\mathrm{d}x = s^* = O(\sqrt{\varepsilon}\,\omega(\varepsilon)),\end{aligned}$$

which yields the desired statement (3.30).

Let us now consider the estimate 3.31. Note by Remark 3.15 the matrix $\tilde{\Sigma}_\tau^{-1}$ given by (3.26) is positive definite and the matrix we subtract is also positive definite. Therefore, it holds in the sense of quadratic forms

$$0 < \tilde{\Sigma}_\tau^{-1} = \Sigma_\tau^{-1} - \frac{1}{\Sigma_\tau^{-1}[\dot{\gamma}_\tau]}\Sigma_\tau^{-1} \otimes \Sigma_\tau^{-1} \le \Sigma_\tau^{-1}.$$

Now, the uniform bound (3.18) yields

$$\sqrt{\det\nolimits_{1,1}(Q_\tau^\top \tilde{\Sigma}_\tau^{-1} Q_\tau)} \le C_\Sigma^{\frac{n-1}{2}}.$$

Then, the desired statement (3.31) follows directly from the estimate (3.25). ☐

3.3.3. Splitting of transport cost and reduction to neighborhood around the saddle

Firstly, observe that from (3.31) follows the a priori estimate

$$\frac{\mathcal{A}^2(x)}{\mu(x)} \lesssim \left(\frac{C_\Sigma}{2\pi\varepsilon}\right)^{n-1} Z_\mu \, e^{\frac{1}{\varepsilon}H(x)} \left(1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))\right). \tag{3.33}$$

Hence, on an exponential scale, the leading order contribution to the cost comes from neighborhoods of points where $H(x)$ is large. Therefore we want to make the set, where H is comparable to its value at the optimal connecting saddle $s_{i,j}$, as small as possible. For this purpose, let us define the following set

$$\Xi_{\gamma,\Sigma} := \left\{ x \in \operatorname{supp}\mathcal{A} : H(x) \geq H(s_{i,j}) - \varepsilon\omega^2(\varepsilon) \right\}. \tag{3.34}$$

Therewith, we obtain by denoting the complement $\Xi^c_{\gamma,\Sigma} := \operatorname{supp}\mathcal{A}\backslash\Xi_{\gamma,\Sigma}$ the splitting

$$\mathcal{T}^2_\mu(\nu_i,\nu_j) \leq \int_{\Xi_{\gamma,\Sigma}} \frac{\mathcal{A}^2(x)}{\mu(x)}\,\mathrm{d}x + \int_{\Xi^c_{\gamma,\Sigma}} \frac{\mathcal{A}^2(x)}{\mu(x)}\,\mathrm{d}x.$$

The integral on $\Xi^c_{\gamma,\Sigma}$ can be estimated with the a priori estimate (3.33) and Lemma 3.16 as follows

$$\begin{aligned}
\int_{\Xi^c_{\gamma,\Sigma}} \frac{\mathcal{A}^2(x)}{\mu(x)}\,\mathrm{d}x &\overset{(3.34)}{\leq} Z_\mu e^{\frac{H(s_{i,j})}{\varepsilon} - \omega^2(\varepsilon)} \int_{\Xi^c_{\gamma,\Sigma}} \mathcal{A}^2(x)\,\mathrm{d}x \\
&\overset{(3.31)}{\lesssim} Z_\mu e^{\frac{H(s_{i,j})}{\varepsilon} - \omega^2(\varepsilon)} \left(\frac{C_\Sigma}{2\pi\varepsilon}\right)^{\frac{n-1}{2}} \int \mathcal{A}(x)\,\mathrm{d}x \\
&\overset{(3.30)}{\lesssim} Z_\mu e^{\frac{H(s_{i,j})}{\varepsilon} - \omega^2(\varepsilon)} \left(\frac{C_\Sigma}{2\pi\varepsilon}\right)^{\frac{n-1}{2}} T.
\end{aligned}$$

Notice that the last estimate yields the second summand in the desired estimate (3.19).

3.3.4. Cost estimate around the saddle

The aim of this subsection is to deduce the estimate

$$\int_{\Xi_{\gamma,\Sigma}} \frac{\mathcal{A}^2(x)}{\mu(x)}\,\mathrm{d}x \lesssim \frac{Z_\mu}{(2\pi\varepsilon)^{\frac{n}{2}}}\, e^{\frac{H(s_{i,j})}{\varepsilon}}\, \frac{2\pi\varepsilon\sqrt{|\det(\nabla^2 H(s_{i,j})|}}{|\lambda^-|}. \tag{3.35}$$

Note that this estimate would yield the missing ingredient for the verification of the desired estimate (3.19).

By the non-degeneracy Assumption 1.3, we can assume that ε is small enough such that $E_0^- \cup E_T^+ \subset \Xi^c_{\gamma,\Sigma}$. It follows that $\Xi_{\gamma,\Sigma} \subset E$. We claim that the transport interpolation Φ_s can be chosen such that there exists a connected interval $I_T \subset [0,T]$ satisfying

$$\Xi_{\gamma,\Sigma} \subset \bigcup_{s\in I_T} (V_s + \gamma_s) \qquad \text{and} \qquad \mathcal{H}^1(I_T) = O(\sqrt{\varepsilon}\,\omega(\varepsilon)). \tag{3.36}$$



Indeed, the level set $\{x \in \mathbb{R}^n : H(x) \le H(s_{i,j}) - \varepsilon\omega^2(\varepsilon)\}$ consists of at least two connected components M_i and M_j such that $m_i \in M_i$ and $m_j \in M_j$. Further, it holds

$$\operatorname{dist}(M_i, M_j) = \inf_{x\in M_i, y\in M_j} |x-y| = O(\sqrt{\varepsilon}\,\omega(\varepsilon)),$$

which follows from expanding H around $s_{i,j}$ in direction of the eigenvector corresponding to the negative eigenvalue of $\nabla^2 H(s_{i,j})$. We can choose the path γ in direction of this eigenvector in a neighborhood of size $O(\sqrt{\varepsilon}\;\omega(\varepsilon))$ around $s_{i,j}$, which shows (3.36).

Combining the covering (3.36) and Lemma 3.13 yields the estimate

$$\int_{\Xi_{\gamma,\Sigma}} \frac{\mathcal{A}^2(x)}{\mu(x)}\,\mathrm{d}x \le \int_{I_T}\int_{V_s} \frac{\mathcal{A}^2(\gamma_s+z_s)}{\mu(\gamma_s+z_s)}\,\mathrm{d}z_s\,\mathrm{d}s. \tag{3.37}$$

Recalling the definition (3.21) of the family of rotations $(Q_\tau)_{\tau\in[0,T]}$, it holds that $z_\tau = Q_\tau z$ with $z \in \{0\}\times\mathbb{R}^{n-1}$. Hence, the following relation holds

$$\int_{I_T}\int_{V_\tau} \frac{\mathcal{A}^2(\gamma_\tau+z_\tau)}{\mu(\gamma_\tau+z_\tau)}\,\mathrm{d}z_\tau\,\mathrm{d}\tau = \int_{\{0\}\times\mathbb{R}^{n-1}}\int_{I_T} \mathbb{1}_{V_\tau}(Q_\tau z)\frac{\mathcal{A}^2(\gamma_\tau+Q_\tau z)}{\mu(\gamma_\tau+Q_\tau z)}\,\mathrm{d}\tau\,\mathrm{d}z. \tag{3.38}$$

The next step is to rewrite $H(\gamma_\tau+Q_\tau z)$. By the reason that $|z_\tau| = O(\sqrt{\varepsilon}\,\omega(\varepsilon))$ for $z_\tau \in V_\tau$ and the global non-degeneracy assumption (1.5), we can Taylor expand $H(\gamma_\tau+z_\tau)$ around $s_{i,j} = \gamma_{\tau^*}$ for $\tau \in I_T$ and $z_\tau = Q_\tau z \in V_\tau$. More precisely, we get

$$\begin{aligned} H(\gamma_\tau+Q_\tau z) - H(s_{i,j}) &= \frac{1}{2}\nabla^2 H(s_{i,j})[\gamma_\tau + Q_\tau z - s_{i,j}] + O(|\gamma_\tau + Q_\tau z - s_{i,j}|^3) \\ &= \frac{1}{2}\nabla^2 H(s_{i,j})[\gamma_\tau - \gamma_{\tau^*}] + \frac{1}{2}\nabla^2 H(s_{i,j})[Q_\tau z] \\ &\quad + \big\langle Q_\tau z, \nabla^2 H(s_{i,j})(\gamma_\tau - \gamma_{\tau^*})\big\rangle + O(|\gamma_\tau + Q_\tau z - \gamma_{\tau^*}|^3) \end{aligned}$$

Now, further expanding γ_τ and Q_τ in τ leads to

$$\gamma_\tau = \gamma_{\tau^*} + \dot\gamma_{\tau^*}(\tau-\tau^*) + O(|\tau-\tau^*|), \quad \text{and} \quad Q_\tau z = Q_{\tau^*}z + O(|\tau-\tau^*|\;|z|).$$

For the expansion of H, we arrive at the identity

$$\begin{aligned} &H(\gamma_\tau+Q_\tau z) - H(s_{i,j}) = \\ &\qquad \frac{1}{2}\nabla^2 H(s_{i,j})[\dot\gamma_{\tau^*}(\tau-\tau^*) + O(|\tau-\tau^*|^2)] + \frac{1}{2}\nabla^2 H(s_{i,j})[Q_{\tau^*}z + O(|\tau-\tau^*|\,|z|)] \\ &\qquad + \Big\langle Q_{\tau^*}z + O(|\tau-\tau^*|\,|z|), \nabla^2 H(s_{i,j})\Big(\dot\gamma_{\tau^*}(\tau-\tau^*) + O(|\tau-\tau^*|^2)\Big)\Big\rangle \\ &\qquad + O(|\gamma_\tau + Q_\tau z - \gamma_{\tau^*}|^3) \\ &= \frac{1}{2}\nabla^2 H(s_{i,j})[\dot\gamma_{\tau^*}]\,(\tau-\tau^*)^2 + \frac{1}{2}\nabla^2 H(s_{i,j})[Q_{\tau^*}z] + \big\langle Q_{\tau^*}z, \nabla^2 H(s_{i,j})\dot\gamma_{\tau^*}\big\rangle\,(\tau-\tau^*) \\ &\qquad + O(|\tau-\tau^*|^3, |z|\,|\tau-\tau^*|^2, |z|^2\,|\tau-\tau^*|, |z|^3). \end{aligned}$$

Using $|\tau - \tau^*| = O(\sqrt{\varepsilon}\ \omega(\varepsilon))$ and $|z| = O(\sqrt{\varepsilon}\ \omega(\varepsilon))$ we obtain for the error the order estimate

$$O(|\tau - \tau^*|^3, |z|\,|\tau - \tau^*|^2, |z|^2\,|\tau - \tau^*|, |z|^3) = O(\varepsilon^{\frac{3}{2}}\ \omega^3(\varepsilon)).$$

The term $\left\langle Q_{\tau^*} z, \nabla^2 H(s_{i,j}) \dot{\gamma}_{\tau^*} \right\rangle (\tau - \tau^*)$ in the expansion of H has no sign and therefore has to vanish. This is only the case, if we choose $\dot{\gamma}_{\tau^*}$ as an eigenvector of $\nabla^2 H(s_{i,j})$ to the negative eigenvalue λ^-, because then

$$\left\langle Q_{\tau^*} z, \nabla^2 H(s_{i,j}) \dot{\gamma}_{\tau^*} \right\rangle (\tau - \tau^*) = \lambda^- \left\langle Q_{\tau^*} z, \dot{\gamma}_{\tau^*} \right\rangle = 0.$$

Additionally, by this choice of $\dot{\gamma}_{\tau^*}$ the quadratic form $\nabla^2 H(s_{i,j})[\dot{\gamma}_{\tau^*}]$ evaluates to

$$\nabla^2 H(s_{i,j})[\dot{\gamma}_{\tau^*}] = \lambda^- |\dot{\gamma}_{\tau^*}|^2 = \lambda^-.$$

Therefore we deduced the desired rewriting of $H(\gamma_\tau + Q_\tau z)$ as

$$H(\gamma_\tau + Q_\tau z) = H(s_{i,j}) - |\lambda^-| (\tau - \tau^*)^2 + \frac{1}{2} \nabla^2 H(s_{i,j})[Q_{\tau^*} z] + O(\varepsilon^{\frac{3}{2}}\ \omega^3(\varepsilon)).$$

From the regularity assumptions on the transport interpolation we can deduce that

$$\begin{aligned} \tilde{\Sigma}_\tau^{-1}[Q_\tau z] &= \tilde{\Sigma}_{\tau^*}^{-1}[Q_\tau z] + O(|\tau - \tau^*|\,|z|^2) \\ &= \tilde{\Sigma}_{\tau^*}^{-1}[Q_{\tau^*} z + O(|\tau - \tau^*|\,|z|)] + O(|\tau - \tau^*|\,|z|^2) \\ &= \tilde{\Sigma}_{\tau^*}^{-1}[Q_{\tau^*} z] + O(\varepsilon^{\frac{3}{2}} \omega^3(\varepsilon)). \end{aligned}$$

Then, it follows easily from the definition (3.25) of P_τ that

$$P_\tau \approx P_{\tau^*}. \tag{3.39}$$

Applying the cost estimate (3.25) of Lemma 3.14, the rewriting 3.3.4 of $H(\gamma_\tau + Q_\tau z)$, and the identity (3.39) yields the estimate

$$\frac{\mathcal{A}^2(\gamma_\tau + Q_\tau z)}{\mu(\gamma_\tau + Q_\tau z)} \lesssim Z_\mu e^{\frac{H(s_{i,j})}{\varepsilon}} P_{\tau^*}^2 e^{-\frac{(2\tilde{\Sigma}_{\tau^*}^{-1} - \nabla^2 H(s_{i,j}))[Q_{\tau^*} z]}{2\varepsilon} - \frac{|\lambda^-|(\tau - \tau^*)^2}{2\varepsilon}}. \tag{3.40}$$

The exponentials are densities of two Gaussian, if we put an additional constraint on the transport interpolation. Namely, we postulate

$$2\tilde{\Sigma}_{\tau^*}^{-1} - \nabla^2 H(s_{i,j}) > 0 \quad \text{on} \quad \operatorname{span} V_{\tau^*}$$

in the sense of quadratic forms. Note that $\operatorname{span} V_{\tau^*} = Q_{\tau^*}(\{0\} \times \mathbb{R}^{n-1}) = \operatorname{span}\{\dot{\gamma}_{\tau^*}\}^\perp$ is the tangent space of the stable manifold in the 1-saddle $s_{i,j}$. With this preliminary considerations we finally are able to estimate the right hand side of (3.38) as

$$\begin{aligned}
&\int_{\{0\}\times\mathbb{R}^{n-1}} \int_{I_T} \mathbb{1}_{V_\tau}(Q_\tau z)\, \frac{\mathcal{A}^2(\gamma_\tau + Q_\tau z)}{\mu(\gamma_\tau + Q_\tau z)} \,\mathrm{d}\tau\, \mathrm{d}z \\
&\overset{(3.40)}{\lesssim} Z_\mu\, e^{\frac{H(s_{i,j})}{\varepsilon}} \int_{\{0\}\times\mathbb{R}^{n-1}} \int_{I_T} P_{\tau^*}^2\, e^{-\frac{(2\tilde{\Sigma}_{\tau^*}^{-1} - \nabla^2 H(s_{i,j}))[Q_{\tau^*} z]}{2\varepsilon} - \frac{\left|\lambda^-\right|(\tau-\tau^*)^2}{2\varepsilon}} \,\mathrm{d}\tau\, \mathrm{d}z \\
&\le Z_\mu\, e^{\frac{H(s_{i,j})}{\varepsilon}}\, \frac{\sqrt{2\pi\varepsilon}}{\sqrt{|\lambda^-|}} \int_{\{0\}\times\mathbb{R}^{n-1}} P_{\tau^*}^2\, e^{-\frac{(2\tilde{\Sigma}_{\tau^*}^{-1} - \nabla^2 H(s_{i,j}))[Q_{\tau^*} z]}{2\varepsilon}} \,\mathrm{d}z \\
&= Z_\mu\, e^{\frac{H(s_{i,j})}{\varepsilon}}\, \frac{\sqrt{2\pi\varepsilon}}{\sqrt{\lambda^-}}\, P_{\tau^*}^2\, \frac{(2\pi\varepsilon)^{\frac{n-1}{2}}}{\sqrt{\det_{1,1}\left(Q_{\tau^*}^\top (2\tilde{\Sigma}_{\tau^*}^{-1} - \nabla^2 H(s_{i,j}))Q_{\tau^*}\right)}} \qquad (3.41)\\
&= \frac{Z_\mu}{(2\pi\varepsilon)^{\frac{n}{2}}}\, e^{\frac{H(s_{i,j})}{\varepsilon}}\, \frac{2\pi\varepsilon}{\sqrt{|\lambda^-|}}\, \underbrace{\frac{\det_{1,1}(Q_{\tau^*}^\top \tilde{\Sigma}_{\tau^*}^{-1} Q_{\tau^*})}{\sqrt{\det_{1,1}\left(Q_{\tau^*}^\top (2\tilde{\Sigma}_{\tau^*}^{-1} - \nabla^2 H(s_{i,j}))Q_{\tau^*}\right)}}}_{\text{to optimize!}} .
\end{aligned}$$

The final step consists of optimizing the choice of $\tilde{\Sigma}_{\tau^*}$. Let us use the notation $A = Q_{\tau^*}^\top \tilde{\Sigma}_{\tau^*}^{-1} Q_{\tau^*}$ and $B = Q_{\tau^*}^\top H(s_{i,j}) Q_{\tau^*}$. Then the minimization problem has the structure

$$\inf_{A\in S^+(n)} \left\{ \frac{\det_{1,1} A}{\sqrt{\det_{1,1}(2A - B)}} : 2A - B > 0 \quad \text{on} \quad \{0\}\times\mathbb{R}^{n-1} \right\}. \qquad (3.42)$$

In the appendix, we show in Lemma B.2 that the optimal value of (3.42) is attained at $\tilde{\Sigma}_{\tau^*}^{-1} = \nabla^2 H(s_{i,j})$ restricted V_{τ^*}. The optimal value is given by

$$\frac{\det_{1,1} A}{\sqrt{\det_{1,1}(2A - B)}} = \sqrt{\det_{1,1}(Q_{\tau^*}^\top \nabla^2 H(s_{i,j}) Q_{\tau^*})}.$$

Because V_{τ^*} is the tangent space of the stable manifold of the saddle $s_{i,j}$ it holds

$$\det_{1,1}(Q_{\tau^*}^\top \nabla^2 H(s_{i,j}) Q_{\tau^*}^\top) = \frac{\det(\nabla^2 H(s_{i,j}))}{\lambda^-} = \frac{|\det(\nabla^2 H(s_{i,j}))|}{|\lambda^-|}. \qquad (3.43)$$

The final step is a combination of (3.37), (3.38), (3.41) and (3.43) to obtain the desired estimate (3.35).

For the verification of Lemma 3.11, it is only left to deduce the estimate (3.20). For that purpose we analyze the error terms in the estimate (3.19) i.e.

$$\mathcal{T}_\mu^2(\nu_i,\nu_j) \lesssim \frac{Z_\mu}{(2\pi\varepsilon)^{\frac{n}{2}}}\, e^{\frac{H(s_{i,j})}{\varepsilon}}\, 2\pi\varepsilon \left(\underbrace{\frac{\sqrt{|\det(\nabla^2 H(s_{i,j})|}}{|\lambda^-|}}_{=O(1)} + \underbrace{\frac{T(C_\Sigma)^{\frac{n-1}{2}}}{\sqrt{2\pi\varepsilon}} e^{-\omega^2(\varepsilon)}}_{=O(\varepsilon^{-\frac{1}{2}} e^{-\omega^2(\varepsilon)})} \right).$$

By our choice of $\omega(\varepsilon) \geq |\ln \varepsilon|^{\frac{1}{2}}$, enforced by Lemma 3.6, we see that

$$O(\varepsilon^{-\frac{1}{2}} e^{-\omega^2(\varepsilon)}) = O(\sqrt{\varepsilon}).$$

Recalling, that "$\lesssim$" means "$\leq$" up to a multiplicative error of order $1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))$ we get

$$\mathcal{T}_\mu^2(\nu_i, \nu_j) \leq \frac{Z_\mu}{(2\pi\varepsilon)^{\frac{n}{2}}}\, e^{\frac{H(s_{i,j})}{\varepsilon}}\, 2\pi\varepsilon \frac{\sqrt{|\det(\nabla^2 H(s_{i,j})|}}{|\lambda^-|} \left(1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))\right) \left(1 + O(\sqrt{\varepsilon})\right).$$

The last inequality already yields the desired estimate (3.20) by using the observation

$$\left(1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))\right) \left(1 + O(\sqrt{\varepsilon})\right) = \left(1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))\right).$$

Remark 3.17. Let us summarize the additional constraints on the transport interpolation besides the Assumption 3.10 of a regular affine transport interpolation to obtain the desired estimate (3.19):

- γ passes the saddle point $s_{i,j}$ at the passage time τ^* in direction of the eigenvector to the negative eigenvalue λ^- of $\nabla^2 H(s_{i,j})$
- γ stays in the sublevel set $\left\{H(x) \leq H(s_{i,j}) - \varepsilon\omega^2(\varepsilon)\right\}$ up to a small time interval of order $\sqrt{\varepsilon}\,\omega(\varepsilon)$ around the passage time τ^*
- It holds $\Sigma_{\tau^*}^{-1} = \nabla^2 H(s_{i,j})$ on the stable manifold of $s_{i,j}$.

3.4. Conclusion of the proof of Theorem 1.9

With the help of Lemma 3.6 and Lemma 3.11 the proof of Theorem 1.9 is straightforward. We can estimate the mean-differences w.r.t. to the measure μ_i by introducing the means w.r.t. the approximations ν_i and ν_j.

$$\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2 = \left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\nu_i}(f) + \mathbb{E}_{\nu_i}(f) - \mathbb{E}_{\nu_j}(f) + \mathbb{E}_{\nu_j}(f) - \mathbb{E}_{\mu_j}(f)\right)^2$$

We apply the Young inequality with a weight that is motivated by the final total multiplicative error term $R(\varepsilon)$ in Theorem 1.9. More precisely,

$$\begin{aligned}\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2 &\leq (1 + \varepsilon^{\frac{1}{2}}\omega^3(\varepsilon)) \left(\mathbb{E}_{\nu_i}(f) - \mathbb{E}_{\nu_j}(f)\right)^2 \\ &\quad + 2(1 + \varepsilon^{-\frac{1}{2}}\omega^{-3}(\varepsilon)) \left(\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\nu_i}(f)\right)^2 + \left(\mathbb{E}_{\mu_j}(f) - \mathbb{E}_{\nu_j}(f)\right)^2\right).\end{aligned}$$

Then, the estimate (3.8) of Lemma 3.6 yields

$$\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2 \leq (1 + \sqrt{\varepsilon}\,\omega^3(\varepsilon)) \left(\mathbb{E}_{\nu_i}(f) - \mathbb{E}_{\nu_j}(f)\right)^2 + O(\varepsilon) \int |\nabla f|^2 \,\mathrm{d}\mu, \tag{3.44}$$

which justifies the statement, that the approximation only leads to higher-order error terms in ε. An application of (3.1) to the estimate (3.44) transfers the mean-difference to the Dirichlet form with the help of the weighted transport distance

$$\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2 \leq \left(\left(1+\sqrt{\varepsilon}\,\omega^3(\varepsilon)\right)\, \mathcal{T}_\mu^2(\nu_i,\nu_j) + O(\varepsilon)\right) \int |\nabla f|^2 \,\mathrm{d}\mu,$$

Since the weighted transport distance $\mathcal{T}_\mu(\nu_i,\nu_j)$ is dominating and we can assume w.l.o.g. that $\mathcal{T}_\mu(\nu_i,\nu_j) \geq 1$, we arrive at the estimate

$$\left(\mathbb{E}_{\mu_i}(f) - \mathbb{E}_{\mu_j}(f)\right)^2 \lesssim \mathcal{T}_\mu^2(\nu_i,\nu_j) \int |\nabla f|^2 \,\mathrm{d}\mu.$$

Now, the Theorem 1.9 follows directly from an application of the estimate (3.20) of Lemma 3.11 and setting $\omega(\varepsilon) = |\ln \varepsilon|^{\frac{1}{2}}$.

4. The local Poincaré inequality: proof of Theorem 1.7

In this section, we are considering only one of the basins of attraction Ω_i and therefore we omit the index i. We will write Ω and μ instead of Ω_i and μ_i respectively. Further, we assume w.l.o.g. that $0 \in \Omega$ is the unique minimum in Ω.

We start with some heuristics for the validity of Theorem 1.7. We consider the Gibbs measure μ restricted to a basin of attraction Ω given by

$$\mu(\mathrm{d}x) = \frac{\mathbb{1}_\Omega(x)}{Z_\mu} \exp\left(-\frac{H(x)}{\varepsilon}\right) \mathrm{d}x.$$

We have to show that this restricted Gibbs measure μ satisfies the PI(ϱ) with constant $\varrho^{-1} = O(\varepsilon)$ (cf. Lemma (1.5)). We argue as follows: On a basin of attraction Ω can only be one local minimum of H, which is located w.l.o.g. at 0. Therefore, on Ω cannot exist another metastable state of the diffusion ξ_t given by (1.3). This means that for small noise ε the diffusion ξ_t slides down on the energy landscape without any obstacle until it reaches a small region around the local minimum of H at 0. This heuristically implies that only the small region around 0 is important for the Poincaré constant ofμ. Therefore, the Gibbs measure μ restricted to the basin of attraction Ω and the Gibbs measure μ restricted to a small neighborhood of 0 should have the same Poincaré constant in terms of scaling in ε. However, around the local minimum 0 the Hamiltonian H is strictly convex due to our non-degeneracy assumption (1.5). Now, an application of the criterion of Bakry-Émery [BÉ85] – it connects convexity of $\frac{H}{\varepsilon}$ to the Poincaré constant of μ – yields that the Gibbs measure μ restricted to a small region around 0 should satisfy the PI(ϱ) with constant $\varrho^{-1} = O(\varepsilon)$.

Now, let us turn to the main idea of our proof of Theorem 1.7. First of all, we want to note that the standard criteria for the Poincaré inequality do not cover our situation.

- The criterion of Bakry-Émery [BÉ85] cannot be applied because H is not convex on the basin of attraction Ω.
- The perturbation lemma of Holley-Stroock [HS87] cannot be applied naively because it would yield an exponentially bad dependence of the Poincaré constant ϱ on ε.

Staying close to the heuristics from above, the Lyapunov condition would be the most natural approach to Theorem 1.7 (cf. Section 4.1 or [BBCG08]). However, one would have to construct an explicit Lyapunov function (cf. Definition 4.4) with rather subtle properties. Especially around critical points x, i.e. $\nabla H(x) = 0$, this seems to be very difficult.

For that reason we choose a different approach. The motivation for our argument comes from the one-dimensional case i.e. $H : \mathbb{R} \to \mathbb{R}$. In this situation, Theorem 1.7 can be deduced by using the Muckenhoupt functional (cf. Section 4.3 and [Muc72]), which determines the Poincaré constant in one dimension up to a universal factor. Because of the non-existence of a multidimensional version of the Muckenhoupt functional, we have to reduce the multi-dimensional case to the one-dimensional case.

For this purpose, we introduce on Ω polar like coordinates $\psi_\eta(r) \in \Omega$ for $\eta \in S^{n-1}$ and $r \geq 0$ (cf. Section 4.2). The coordinates $\psi_\eta(r)$ are going to be a small perturbation of the coordinates resulting from the deterministic gradient-flow

$$\frac{\mathrm{d}}{\mathrm{d}t}\xi_t = -\nabla H(\xi_t),$$

parameterized by arc-length i.e. $|\dot{\psi}_\eta(r)| = 1$. In these coordinates, the restricted Gibbs measures becomes

$$\mu(\mathrm{d}r, \mathrm{d}\eta) = \frac{\mathbb{1}_\Omega(\psi_\eta(r))}{Z}\ j_\eta(t)\ \exp\left(-\frac{1}{\varepsilon}H(\psi_\eta(r))\right)\ \mathrm{d}r\ \mathrm{d}\eta, \tag{4.1}$$

where $j_\eta(r)$ is the Jacobian determinant of the coordinate transformation.

Now, we carry out a two-scale argument similar to the one used for the proof of the Eyring-Kramers formula (see Corollary 1.10 and Section 2). So, the restricted Gibbs measure $\mu(\mathrm{d}r, \mathrm{d}\eta)$ is decomposed into

$$\mu(\mathrm{d}r, \mathrm{d}\eta) = \mu(\mathrm{d}r|\eta)\ \hat{\mu}(\mathrm{d}\eta), \tag{4.2}$$

where the conditional measures $\mu(\mathrm{d}r|\eta)$ and the marginal $\hat{\mu}(\mathrm{d}\eta)$ are given by

$$\mu(\mathrm{d}r|\eta) = \frac{1}{\hat{\mu}(\eta)} e^{-\frac{1}{\varepsilon}H(\psi_\eta(r))} j_\eta(r)\ \mathrm{d}r \tag{4.3}$$

$$\hat{\mu}(\mathrm{d}\eta) = \frac{1}{Z_\mu}\int e^{-\frac{1}{\varepsilon}H(\psi(r))} j_\eta(r)\ \mathrm{d}r\ \mathrm{d}\eta. \tag{4.4}$$

Therewith, we get the same splitting as in the discrete case of Lemma 1.6, i.e.

$$\begin{aligned}\operatorname{var}_{\mu(\mathrm{d}x)}(f(x)) &= \operatorname{var}_{\mu(\mathrm{d}r,\mathrm{d}\eta)}(f(\psi_\eta(r))) \\ &= \mathbb{E}_{\hat\mu(\mathrm{d}\eta)}\left(\operatorname{var}_{\mu(\mathrm{d}r|\eta)}(f(\psi_\eta(r)))\right) + \operatorname{var}_{\hat\mu(\mathrm{d}\eta)}\left(\mathbb{E}_{\mu(\mathrm{d}r|\eta)}(f(\psi_\eta(r)))\right). \quad (4.5)\end{aligned}$$

Let us consider the first term on the right hand side of the last equation. Note that the conditional measures $\mu(\mathrm{d}r|\eta)$ are one-dimensional. Hence, we are able to deduce Poincaré inequalities for the conditional measures $\mu(\mathrm{d}r|\eta)$ using the Muckenhoupt functional. Of course, this step is very sensitive to the choice of the coordinates $\{\psi_\eta\}_{\eta\in S^{n-1}}$. However, a careful and non-trivial construction of the coordinates yields the desired scaling of the Poincaré constant. The next statement contains the existence of *good* coordinates $\{\psi_\eta\}_{\eta\in S^{n-1}}$.

Proposition 4.1. *Assume that the basin of attraction Ω is bounded. Then, there exist coordinates $\{\psi_\eta\}_{\eta\in S^{n-1}}$ on Ω such that the conditional measures $\mu(\mathrm{d}r|\eta)$ are radial-mixing measures in the sense of Definition 4.14.*

The definition of radial-mixing measures is technically and for the moment means that the Hamiltonian evaluated along coordinates lines $H(\psi_\eta(r))$ fulfills certain monotonicity assumptions and the Jacobian determinant $j_\eta(r)$ of the coordinate transform shows a controlled blowup behavior, not causing any metastabilites. Because the proof of Proposition 4.1 is relatively elementary but a bit lengthy, it is carried out in full detail in the Appendix A. We want to point out that Proposition 4.1 demands the boundedness of the domain Ω. However, this does not present a problem, because in Section 4.1 we will show by a Lyapunov argument that we may indeed assume that Ω is bounded (cf. Corollary 4.7 below).

The next statement shows that radial-mixing measures are satisfying the Poincaré inequality $\mathrm{PI}(\varrho)$ with the right-scaling Poincaré constant ϱ.

Proposition 4.2. *Let $\mu(\cdot|\eta)$ be a radial-mixing measure in the sense of Definition 4.14. Then the measure $\mu(\cdot|\eta)$ satisfies $\mathrm{PI}(\varrho(\eta))$ with constant*

$$\frac{1}{\varrho(\eta)} = O(\varepsilon) \qquad \textit{for} \quad \varepsilon\to 0$$

uniformly in $\eta\in S^{n-1}$.

The proof of Proposition 4.2 is carried out in the Section 4.3. It is based on some properties of radial-mixing measures that are outlined in Section 4.2.

Now, let us consider the second term on the right hand side of (4.5), which is given by

$$\operatorname{var}_{\hat\mu(\mathrm{d}\eta)}(\hat f(\eta)) = \int\left(\hat f(\eta) - \int \hat f(\theta)\ \hat\mu(d\theta)\right)^2 \hat\mu(\mathrm{d}\eta) \qquad (4.6)$$

by using the notation

$$\hat{f}(\eta) = \mathbb{E}_{\mu(\mathrm{d}r|\eta)}(f(\psi_\eta(r))).$$

Unfortunately, an application of the same strategy as in [GOVW09] would not yield the desired estimate. The reason is that, following [GOVW09], one would apply a Poincaré inequality for the marginal measure $\hat{\mu}(\mathrm{d}\eta)$ and then estimate the coarse-grained gradient by the full gradient via a covariance estimate. However, the Poincaré constant for the marginal measure $\hat{\mu}$ already is at least of order 1, since $\hat{\mu}$ lives on the sphere S^{n-1}. Therefore, this strategy could not yield the desired Poincaré inequality $\mathrm{PI}(\varrho)$ with constant of order $\varrho^{-1} = O(\varepsilon)$.

For that reason we use a different strategy. Following the proof of Corollary 1.10, we represent the polar variance given by (4.6) as mean-difference. More precisely, the polar variance takes the form

$$\operatorname{var}_{\hat{\mu}(\mathrm{d}\eta)}(\hat{f}(\eta)) = \frac{1}{2} \int \int \left(\hat{f}(\eta) - \hat{f}(\theta) \right)^2 \hat{\mu}(\mathrm{d}\eta)\, \hat{\mu}(\mathrm{d}\theta). \tag{4.7}$$

From this starting point, the procedure is somehow similar to the proof of the mean-difference estimate of Theorem 1.9. However, the procedure needs some more evolved ingredients due to the fact that the marginal measure $\hat{\mu}$ lives on the continuous state space S^{n-1} and the conditional measures $\mu(\mathrm{d}r|\eta)$ have one-dimensional support. The argument is outlined in detail in Section 4.4, in which the following statement is deduced.

Proposition 4.3 (Polar mean-difference estimate)**.** *It holds the estimate*

$$\int \int \left(\hat{f}(\eta) - \hat{f}(\theta) \right)^2 \hat{\mu}(\mathrm{d}\eta)\, \hat{\mu}(\mathrm{d}\theta) \leq O(\varepsilon) \int |\nabla f|^2 \,\mathrm{d}\mu.$$

Now, we have provided all the ingredients that are needed for the proof of Theorem 1.7.

Proof of Theorem 1.7. The starting point of the proof is formula (4.5). By Corollary 4.7 from below we may assume w.l.o.g. that Ω is bounded. Therefore, Proposition 4.1 allows us to apply the local Poincaré inequality of Proposition 4.2, which leads to the estimate

$$\begin{aligned} \operatorname{var}_{\mu(\mathrm{d}x)}(f(x)) &\leq O(\varepsilon) \int \int |\partial_r f(\psi_\eta(r))|^2 \,\mu(\mathrm{d}r|\eta)\, \hat{\mu}(\mathrm{d}\eta) + \operatorname{var}_{\hat{\mu}(\mathrm{d}\eta)}(\hat{f}(\eta)) \\ &= O(\varepsilon) \int \int |\nabla f(\psi_\eta(r))|^2 \underbrace{|\dot{\psi}_\eta(r)|^2}_{=1} \mu(\mathrm{d}r, \mathrm{d}\eta) + \operatorname{var}_{\hat{\mu}(\mathrm{d}\eta)}(\hat{f}(\eta)). \end{aligned}$$

We now need to consider the second term on the right hand side of the last inequality. The polar variance $\operatorname{var}_{\hat{\mu}(\mathrm{d}\eta)}(\hat{f}(\eta))$ is expressed by (4.7) as a mean-difference. Hence, an application of the estimate of Proposition 4.3 yields the desired statement. □

It is only left to verify the ingredient used in the proof of Theorem 1.7. In the Section 4.1, we deduce Corollary 4.7. In Section 4.2, we introduce the notion of radial-mixing measures and state some important properties of them. In the Appendix A, we carry out the proof of Proposition 4.1. In Section 4.3, we will prove Proposition 4.2. Finally, in Section 4.4, we state the proof of Proposition 4.3.

4.1. Reduction to bounded domains

In this subsection, we will show that by condition (**A1**) and (**A2**) the Poincaré constant of μ is dominated by the Poincaré constant of a restricted measure μ_R (cf. Corollary 4.7), which is given for some $R > 0$ by the density

$$\mu_R(\mathrm{d}x) = \frac{\mathbb{1}_{\Omega \cap B_R(0)}(x)}{Z}\ \mu(\mathrm{d}x).$$

For this purpose, we apply a standard technique in the context of Poincaré inequalities: the Lyapunov condition.

Definition 4.4 (Lyapunov condition)**.** Let $H : \Omega \to \mathbb{R}$ be a smooth Hamiltonian and let

$$\mu(\mathrm{d}x) = \frac{\mathbb{1}_{\Omega}(x)}{Z}\ \exp\left(-\frac{H(x)}{\varepsilon}\right) \mathrm{d}x$$

denote the associated Gibbs measure μ. Then, $W : \Omega \to [1, \infty)$ is a Lyapunov function for H provided there exist constants $R > 0$, $b > 0$, and $\lambda > 0$ such that

$$\frac{1}{\varepsilon}\ LW \leq -\lambda W + b\, \mathbb{1}_{\Omega_R}. \tag{4.8}$$

For a good paper about the use of the Lyapunov condition for deducing Poincaré inequalities we recommend [BBCG08]. The main ingredient of this technique is the following statement.

Theorem 4.5 (Theorem 1.4. of [BBCG08])**.** *Suppose that H fulfils the Lyapunov condition (4.8) and that μ_R satisfies $\mathrm{PI}(\varrho_R)$. Then also μ satisfies $\mathrm{PI}(\varrho)$ with constant*

$$\varrho \geq \frac{\lambda \varrho_R}{b + \varrho_R}.$$

Even if the proof of Theorem 4.5 is simple, we refer the reader to [BBCG08]. We want to apply Theorem 4.5 to our situation. Hence, we do not only have to verify the Lyapunov condition (4.8) but also have to investigate the dependence of the constants R, b, and λ on the parameter ε. This is carried out in the next lemma.

Lemma 4.6. *Assume that H satisfies the assumption (**A1**) and (**A2**) with constants $C_H >$ and $K_H \geq 0$. Additionally, assume that*

$$\varepsilon \leq \frac{1}{2} \frac{C_H^2}{C_H^2 + 4K_H}.$$

Then H satisfies the Lyapunov condition (4.8) for some ε-independent constants R and b and an ε-dependent constant $\lambda = \frac{C_H^2}{8\varepsilon}$.

Proof. We take $W = e^H$ as Lyapunov function and observe that

$$\frac{1}{\varepsilon}\frac{Le^H}{e^H} = \Delta H + |\nabla H|^2 - \varepsilon^{-1}|\nabla H|^2 = -\Big((\varepsilon^{-1}-2)|\nabla H|^2 + (|\nabla H|^2 - \Delta H)\Big). \tag{4.9}$$

Now, by the assumptions (**A1**) and (**A2**) we find $R < \infty$ such that

$$\forall |x| \geq R : |\nabla H(x)| \geq \tfrac{1}{2}C_H \quad \text{and} \quad |\nabla H(x)|^2 - \Delta H(x) \geq -2K_H.$$

Therewith, we estimate (4.9) in the case $|x| > R$ as

$$\frac{1}{\varepsilon}\frac{Le^H}{e^H} \leq -\frac{1}{4\varepsilon}C_H^2 + \frac{1}{2}C_H^2 + 2K_H \leq -\frac{1}{8\varepsilon}C_H^2 =: -\lambda, \quad \text{provided that } \varepsilon \leq \frac{1}{2}\frac{C_H^2}{C_H^2 + 4K_H}.$$

In the case $|x| \leq R$, we get from (4.9) by omitting the term with negative sign

$$\frac{1}{\varepsilon}\frac{Le^H}{e^H} \leq \Delta H + |\nabla H|^2 \leq \sup_{|x|\leq R}\Big(\Delta H + |\nabla H|^2\Big) =: b,$$

which completes the argument. □

As a simple consequence of Theorem 4.5 and Lemma 4.6, we obtain the main result of this subsection, which allows to consider w.l.o.g. bounded domains Ω for the proof of Theorem 1.7.

Corollary 4.7. *Assume that H satisfies the assumption (**A1**) and (**A2**). Additionally, assume that μ_R satisfies* PI(ϱ_R) *with constant $\frac{1}{\varrho_R} = O(\varepsilon)$. Then μ also satisfies* PI(ϱ) *with constant $\frac{1}{\varrho} = O(\varepsilon)$.*

Proof. We can apply Theorem 4.5 and with Lemma 4.6 and find

$$\frac{1}{\varrho} \leq \frac{b+\varrho_R}{\lambda\varrho_R} = \frac{8\varepsilon}{C_H^2}\left(\frac{b}{\varrho_R}+1\right) = 8\varepsilon\left(O(\varepsilon)+1\right) = O(\varepsilon).$$

□

4.2. The right coordinates and properties of radial-mixing measures

In this subsection we discuss the right choice of local coordinates $(\psi_\eta)_{\eta\in S^{n-1}}$ in order to apply the two-scale proof of Theorem 1.7 of the last section. Moreover, we introduce the notion of radial-mixing measures (cf. Definition 4.14) and state some properties, which are needed in the proof of Proposition 4.2 in Section 4.3.

An obvious question is: Why are we not using the coordinates $\psi_\eta(r)$ resulting from the deterministic gradient-flow

$$\frac{\mathrm{d}}{\mathrm{d}t}\xi_t = -\nabla H(\xi_t),$$

parameterized by arc-length i.e. $|\dot{\psi}_\eta(r)| = 1$? The Poincaré inequality of μ is strongly connected to ergodic properties of the diffusion ξ_t given by (1.3). Intuitively, these coordinates should describe the diffusion ξ_t very well as we are only interested in the regime of vanishing noise ε. For this reason, deterministic gradient-flow coordinates $\psi_\eta(r)$ should be a natural choice of coordinates.

However, we are not using these deterministic gradient-flow coordinates because the conditional measures

$$\mu(\mathrm{d}r|\eta) = \frac{1}{Z} e^{-\frac{1}{\varepsilon} H(\psi_\eta(r))} j_\eta(r)\ \mathrm{d}r$$

given by (4.3) will not show the right scaling of the Poincaré constant. A detailed analysis reveals that problems only arise in neighborhoods of critical points x, i.e. $\nabla H(x) = 0$. Therefore, we will only change the coordinates in neighborhoods around critical points (cf. (**H̃1**) of Assumption 4.8). Heuristically, this observation seems to be plausible as, at critical points, the gradient of H becomes comparable to the noise. Hence, at least there, stochastic effects have to be taken into account. The main stochastic effect to be considered is that the noise does not have a preferred direction. Thus, it is better to smoothly transform the deterministic gradient-flow coordinates around local extrema to polar coordinates (cf. (**H̃2**) of Assumption 4.8).

However, a more serious problem arises if one considers directions $\eta \in S^{n-1}$ such that the coordinate lines $\psi_\eta(r)$ almost hit a saddle point of H. For such directions η, the Hamiltonian

$$H_\eta(r) := \frac{1}{\varepsilon} H(\psi_\eta(r)) - \ln j_\eta(r) \qquad (4.10)$$

of the measure $\mu(\mathrm{d}r|\eta)$ has a metastability in the case of deterministic gradient-flow coordinates $\psi_\eta(r)$, since $j_\eta(r)$ becomes unbounded to $+\infty$. Here, the following stochastic effect has to be taken into account: If one follows a trajectory of the diffusion ξ_t given by (1.3) starting anywhere above the saddle, one will almost never get too close to the saddle point. The reason is that the process ξ_t will be pushed away from the saddle as soon as the noise becomes comparable to $|\nabla H(\xi_t)|$. This stands in contrast to deterministic gradient-flow coordinates, where you can get as close to the saddle as you want. Heuristically, this shows that deterministic gradient-flow coordinates are not appropriate at the saddle for the stochastic dynamics.



Figure 2: Deterministic gradient-flow coordinates obtained from $\tilde{H}$

As you can see by Figure 2, we adjust the deterministic gradient-flow coordinates around the saddles in such a way that you will never get close to the saddle by following a coordinate line from above the saddle. The origin of this idea goes back to a discussion of the authors with Martin Hairer. Technically, these ideas are manifested in the properties (**$\tilde{\mathbf{H}}$3**), (**$\tilde{\mathbf{H}}$4**), and (**$\tilde{\mathbf{H}}$5**). Later, these properties guarantee in the proof of Proposition 4.2 that in the Hamiltonian $H_\eta(r)$ there are no metastabilites for any direction η (cf. Section 4.3).

As we have outlined above, the desired coordinates $\psi_\eta(r)$ will be slight modifications of the coordinates resulting from the deterministic gradient-flow w.r.t. the Hamiltonian H. An easy way to construct such coordinates is to consider a small perturbation $\tilde{H}$ of the Hamiltonian H. Then, we define the modified coordinates $\psi_\eta(r)$ by the deterministic gradient-flow w.r.t. the perturbation $\tilde{H}$. In the next assumption we state some properties of the perturbation $\tilde{H}$ that guarantee that the resulting coordinates will serve our needs.

Assumption 4.8 (Perturbation $\tilde{H}$ of H)**.** *We assume that the domain Ω is bounded. Additionally, we assume that the perturbation $\tilde{H} \in C^2(\Omega \backslash \{0\}) \cap C^0(\bar{\Omega})$ of the Hamiltonian H satisfies the conditions:*

(i) *$\tilde{H}$ equals H up to small δ-neighborhoods around the critical points of H, i.e. by recalling that $\mathcal{S}$ is the set of critical points of H we have*

$$\forall x \notin \bigcup_{y \in \mathcal{S}} B_\delta(y): \ \tilde{H}(x) = H(x). \tag{$\tilde{\mathbf{H}}$1}$$

(ii) *$\tilde{H}$ restricted to a ball around the local minima located in 0 of radius $\frac{\delta}{2} > 0$ is spherically symmetric and linear, i.e.*

$$\forall x \in B_{\frac{\delta}{2}}(0): \ \tilde{H}(x) = \tilde{H}(0) + |x| . \tag{$\tilde{\mathbf{H}}$2}$$

(iii) *$|\nabla \tilde{H}|$ is uniformly bounded from below in $\Omega \backslash \{0\}$, i.e.*

$$\exists c_{\tilde{H}} > 0 \ \forall x \in \Omega \backslash \{0\}: \ |\nabla \tilde{H}(x)| \geq c_{\tilde{H}}. \tag{$\tilde{\mathbf{H}}$3}$$

By using the Assumption 4.8, we are able to define the coordinates $\psi_\eta(r)$ as follows.

Definition 4.9 (Local coordinates $\psi_\eta(r)$)**.** Assume that $\tilde{H}$ satisfies the Assumption 4.8. For $r \in \mathbb{R}$, we consider the associated flow $\Psi_r : \Omega \backslash \{0\} \to \Omega \backslash \{0\}$ defined as the solution of

$$\Psi_0(x) = x \qquad \text{and} \qquad \dot{\Psi}_r(x) = \tilde{F}(\Psi_r(x)) := \frac{\nabla \tilde{H}(\Psi_r(x))}{|\nabla \tilde{H}(\Psi_r(x))|}. \tag{4.11}$$

For $\eta \in S^{n-1}$, the coordinates $\psi_\eta : (0, T_\eta) \to \mathbb{R}^n$ are defined by

$$\psi_\eta(r) = \Psi_{r - \frac{\delta}{2}}(\tfrac{\delta}{2}\eta), \tag{4.12}$$

where the domain $(0, T_\eta)$ is chosen to be maximal. It easily follows from the Assumption 4.8 that there is a constant $C_{\tilde{H}} < \infty$ such that

$$T_\eta < C_{\tilde{H}} \qquad \text{for all } \eta \in S^{n-1}. \tag{4.13}$$

Remark 4.10. Note that the coordinates ψ_η are well-defined by the property (**$\tilde{\mathbf{H}}$2**). Namely, the coordinates $\psi_\eta(r)$ satisfy for $r \in (0, \frac{\delta}{2})$ the equation

$$\dot{\psi}_\eta(r) = \frac{\nabla \tilde{H}(\psi_\eta(r))}{|\nabla \tilde{H}(\psi_\eta(r))|} = \eta. \tag{4.14}$$

Therefore $\Psi_{r-\frac{\delta}{2}}(\frac{\delta}{2}\eta) = r\eta$ for $r \in (0, \frac{\delta}{2})$. This shows that $\psi_\eta(r)$ are polar coordinates around the local minima.

As we have explained above, it is very important to have a good understanding of the Hamiltonian

$$H_\eta(r) = \frac{1}{\varepsilon} H(\psi_\eta(r)) - \ln j_\eta(r)$$

of the conditional measure $\mu(\mathrm{d}r|\eta)$. In the next statement, we consider the evolution of the second term of the last identity, namely the Jacobian determinant $j_\eta(r)$.

Lemma 4.11 (Evolution of $j_\eta(r)$)**.** *For $x \in \Omega \setminus \{0\}$ let $D\Psi_r(x)$ be the Jacobian of the flow Ψ_r given by (4.11). Then the Jacobian determinant*

$$j_\eta(r) := \ \det D\Psi_{s-\frac{\delta}{2}}(\tfrac{\delta}{2}\eta)$$

evaluated along the gradient-flow lines $\psi_\eta(r)$ (cf. (4.12)) satisfies the evolution

$$\forall r \in (0, T_\eta): \qquad \frac{\mathrm{d}}{\mathrm{d}r} \ln j_\eta(r) = \nabla \cdot \tilde{F}(\psi_\eta(r)). \tag{4.15}$$

Proof. Differentiating the identity (4.11) leads to

$$D\dot{\Psi}_r(x) = D\tilde{F}(\Psi_r(x) D\Psi_r(x).$$

As $D\Psi_r$ is regular, we can multiply by $(D\Psi_r)^{-1}(x)$ and take the trace

$$\operatorname{tr}\left(D\dot{\Psi}_r(x)(D\Psi_r)^{-1}(x)\right) = \operatorname{tr} D\tilde{F}(\Psi_r(x)) = \nabla \cdot \tilde{F}(\Psi_r(x)).$$

By Jacobi's formula (B.3) the left hand side of the last identity can be written as

$$\frac{\mathrm{d}}{\mathrm{d}r} \ln \det D\Psi_r(x) = \operatorname{tr}\left(D\dot{\Psi}_r(x) D\Psi_r^{-1}(x)\right),$$

which yields the desired formula (4.15). □

With the help of the last statement, we are able to formulate a condition (cf. (**$\tilde{\mathbf{H}}$4**) below) on the perturbed Hamiltonian $\tilde{H}$. The condition (**$\tilde{\mathbf{H}}$4**) ensures that the Jacobian determinant $j_\eta(r)$ behaves in the right way when applying the Muckenhoupt functional later (cf. Proposition 4.2).

Lemma 4.12 (Bounds on $j_\eta(r)$)**.** *Assume that the perturbed Hamiltonian $\tilde{H}$ satisfies the Assumption 4.8 and that the local coordinates $\psi_\eta(r)$ are given by Definition 4.9. Additionally, assume that $\tilde{H}$ satisfies the condition for $r \geq \delta/2$*

$$-C_{\tilde{H}} \max\left\{1, (T_\eta - r)^{-2}\right\} \leq \nabla \cdot \tilde{F}(\psi_\eta(r)) \leq C_{\tilde{H}}, \tag{$\tilde{\mathbf{H}}$4}$$

for some constant $C_{\tilde{H}} > 1$. Then the Jacobian determinant $j_\eta(r)$ satisfies

$$\forall r \in (0, \tfrac{\delta}{2}] : \quad j_\eta(r) = r^{n-1}, \tag{j1}$$

$$\forall r \in (\tfrac{\delta}{2}, \tilde{T}_\eta) : \quad c_{\tilde{H}} \min\{1, \exp(-C_{\tilde{H}}(T_\eta - r)^{-1})\} \leq j_\eta(r) \leq C_{\tilde{H}} \tag{j2}$$

for some constants $0 < c_{\tilde{H}} < C_{\tilde{H}} < \infty$, uniformly in η.

Proof of Lemma 4.12. The property (**j1**) is a direct consequence of the fact that the coordinates $\psi_\eta(r)$ are spherical coordinates for $r < \frac{\delta}{2}$ by Remark 4.10.
Let us consider the property (**j2**). By the fundamental theorem of calculus and the identity (4.15) we obtain for $r \in (\frac{\delta}{2}, T_\eta)$

$$\ln j_\eta(r) - \ln j_\eta(\tfrac{\delta}{2}) = \int_{\frac{\delta}{2}}^{r} \nabla \cdot \tilde{F}(\psi_\eta(s)) \mathrm{d}s.$$

Taking the exponential in the last identity leaves us with

$$j_\eta(r) = j_\eta(\tfrac{\delta}{2}) \exp\left(\int_{\frac{\delta}{2}}^{r} \nabla \cdot \tilde{F}(\psi_\eta(s)) \mathrm{d}s \right).$$

Employing the uniform upper bound on $T_\eta < C_{\tilde{H}}$ (cf. (4.13)) and the upper bound of ($\tilde{\mathbf{H}}$**4**) immediately yields the desired upper bound on $j_\eta(r)$.
Let us now consider the lower bound of (**j2**). Straightforward calculation yields

$$\begin{aligned}
\exp\left(\int_{\frac{\delta}{2}}^{r} \nabla \cdot \tilde{F}(\psi_\eta(s)) \mathrm{d}s \right) &\geq \exp\left(-C_{\tilde{H}} \int_{\frac{\delta}{2}}^{r} \max\left\{1, (T_\eta - s)^{-2}\right\} \mathrm{d}s \right) \\
&\geq e^{-C_{\tilde{H}} r} \exp\left(-C_{\tilde{H}} \int_{T_\eta - 1}^{\max\{r, T_\eta - 1\}} (T_\eta - s)^{-1} \mathrm{d}s \right) \\
&= e^{-C_{\tilde{H}} r} \exp\left(-C_{\tilde{H}} (T_\eta - \max\left\{T_\eta - 1, r\right\})^{-1} \right).
\end{aligned}$$

Employing again the uniform upper bound on $T_\eta < C_{\tilde{H}}$ (cf. (4.13)) yields the desired lower bound. □

We still need to understand the first term in the Hamiltonian H_η given by (4.10), namely $H(\psi_\eta(r))$. The growth estimates (**h1**) and (**h2**) are one of the main ingredients to apply of the Muckenhoupt functional in the proof of Proposition 4.2 (see Section 4.3).

Lemma 4.13 (Properties of $h_\eta(\cdot)$). *Assume that the perturbed Hamiltonian $\tilde{H}$ satisfies the Assumption 4.8 and that the local coordinates $\psi_\eta(r)$ are given by Definition 4.9. Additionally, assume that $\tilde{H}$ satisfies the condition*

$$\langle \nabla H(\psi_\eta(r)), \nabla \tilde{H}(\psi_\eta(r)) \rangle \geq c_{\tilde{H}} \min\{1, r, T_\eta - r\} \, |\nabla \tilde{H}(\psi_\eta(r))| \tag{$\mathbf{\tilde{H}5}$}$$

for some constant $0 < c_{\tilde{H}}$.
Then, the following estimates hold uniformly in η:

$$\exists \lambda_1 > 0 \ \forall s \in [0, T_\eta) : H(\psi_\eta(r)) \geq H(\psi_\eta(s)) + \lambda_1 (r-s)^2, \quad \forall r \in [s, T_\eta] \tag{$\mathbf{h1}$}$$

$$\exists \lambda_2 > 0 \ \forall s \in (0, T_\eta] : H(\psi_\eta(r)) \leq H(\psi_\eta(sr)) - \lambda_2 (r-s)^2, \quad \forall r \in [0, s]. \tag{$\mathbf{h2}$}$$

Proof of Lemma 4.13. For $r \leq T_\eta$ it follows from Definition 4.9 and ($\mathbf{\tilde{H}5}$) that

$$\frac{\mathrm{d}}{\mathrm{d}r} H(\psi_\eta(r)) = \left\langle \nabla H(\psi_\eta(r)), \frac{\nabla \tilde{H}(\psi_\eta(r))}{|\nabla \tilde{H}(\psi_\eta(r))|} \right\rangle \geq c_{\tilde{H}} \min\{1, r, T_\eta - r\}.$$

Hence on $(0, T_\eta)$, the radial Hamiltonian $h_\eta(r)$ is strictly increasing. Additionally, we have with (4.13) that $T_\eta \leq C_{\tilde{H}} < \infty$ holds uniformly in η. Therefore, it is easy to deduce ($\mathbf{h1}$) and ($\mathbf{h2}$) with constants λ_1 and λ_2 uniformly in η. □

Now, we have provided all the ingredients needed for the proof of Proposition 4.2 in Section 4.3. For convenience, we make the following definition.

Definition 4.14 (Radial-mixing measures). Assume that the perturbed Hamiltonian $\tilde{H}$ satisfies ($\mathbf{\tilde{H}1}$), ($\mathbf{\tilde{H}2}$) and ($\mathbf{\tilde{H}3}$) of Assumption 4.8 and that the local coordinates $\psi_\eta(r)$ are given by Definition 4.9. Additionally, assume that $\tilde{H}$ satisfies the conditions ($\mathbf{\tilde{H}4}$) and ($\mathbf{\tilde{H}5}$). Then, the conditional measures

$$\mu(\mathrm{d}r|\eta) = j_\eta(r) \exp\left(-\frac{1}{\varepsilon} H(\psi_\eta(r))\right) \mathrm{d}r$$

are called *radial-mixing measures.*

The existence of radial-mixing measures is shown in Proposition 4.1, which is verified in the Appendix A. The term *mixing* in the definition of radial-mixing measures is justified by Proposition 4.2, which proofs that the Poincaré constant $\varrho(\eta)$ of $\mu(\mathrm{d}r|\eta)$ satisfies $\varrho^{-1}(\eta) = O(\varepsilon)$.

4.3. The Muckenhoupt functional: proof of Proposition 4.2

The aim of this section is the proof of Proposition 4.2. In the argument we apply the Muckenhoupt functional [Muc72]. For the convenience of the reader, we restate the main result of [Muc72] in Theorem 4.15 below and clarify the connection of the Muckenhoupt functional to the Poincaré inequality in Proposition 4.16. For the sake of completeness we also state the proofs of Theorem 4.15 and Proposition 4.16 in full detail.

The following theorem is a weighted Hardy inequality stated by [Muc72] in a more general context with a sharp characterization of the optimal constant up to a factor 4.

Theorem 4.15 ([Muc72, Theorem 4])**.** *Let μ be an absolutely continuous measure on $\mathbb{R}^+$ with values in $\mathbb{R}^+ \cup \{+\infty\}$, then there exists a constant $C < \infty$ for which*

$$\forall f : \mathbb{R}^+ \to \mathbb{R} \qquad \int_0^\infty \left(\int_0^x f(t)\, \mathrm{d}t \right)^2 \mu(\mathrm{d}x) \le C \int_0^\infty f^2 \mu(\mathrm{d}x), \tag{4.16}$$

if and only if

$$B = \sup_{r>0} \int_r^\infty \mathrm{d}\mu \int_0^r \frac{1}{\mu(x)}\, \mathrm{d}x < \infty. \tag{4.17}$$

Furthermore, the optimal C obeys the estimate

$$B \le C \le 4B. \tag{4.18}$$

Proof of Theorem 4.15. The proof consists of two steps. In the first step we show that (4.17) implies (4.16) with constant $C \le 4B$. In the second step, we will show that (4.16) implies (4.17) by the the estimate $B \le C$, which then establishes (4.18).

Define the function $G(x) = \int_0^x \frac{1}{\mu(y)} \mathrm{d}y$. The first step is to use Cauchy-Schwarz inequality for the inner integral

$$\left(\int_0^x f(t) \mathrm{d}t \right)^2 \le \int_0^x |f(t)|^2 \sqrt{G(t)}\, \mu(\mathrm{d}t) \int_0^x \frac{1}{G(t)\mu(t)}\, \mathrm{d}t \tag{4.19}$$

Now, note that with $2 \frac{\mathrm{d}}{\mathrm{d}t} \sqrt{G(t)} = \frac{1}{\sqrt{G(t)\mu(t)}}$ the last integral evaluates to

$$\int_0^x \frac{1}{G(t)\mu(t)}\, \mathrm{d}t = 2\sqrt{G(x)} \le 2\sqrt{B} \left(\int_x^\infty \mathrm{d}\mu \right)^{-\frac{1}{2}}, \tag{4.20}$$

where the last estimate follows from the definition of the constant B. Combining (4.20) and (4.19), integrating the resulting inequality w.r.t. to μ and using Fubini's theorem to interchange the integration, we arrive at

$$\int_0^\infty \left(\int_0^x f(t)\, \mathrm{d}t \right)^2 \mu(\mathrm{d}x) \le 2\sqrt{B} \int_0^\infty |f(t)|^2 \sqrt{G(t)} \left(\int_t^\infty \left(\int_x^\infty \mathrm{d}\mu \right)^{-\frac{1}{2}} \mu(\mathrm{d}x) \right) \mu(\mathrm{d}t). \tag{4.21}$$

Now, we observe that

$$2 \frac{\mathrm{d}}{\mathrm{d}x} \left(\int_x^\infty \mathrm{d}\mu \right)^{\frac{1}{2}} = - \left(\int_x^\infty \mathrm{d}\mu \right)^{-\frac{1}{2}} \mu(x).$$

Therewith, the inner integrals in (4.21) evaluate to

$$\int_t^\infty \left(\int_x^\infty \mathrm{d}\mu\right)^{-\frac{1}{2}} \mu(\mathrm{d}x) = -2\int_t^\infty \frac{\mathrm{d}}{\mathrm{d}x}\left(\int_x^\infty \mathrm{d}\mu\right)^{\frac{1}{2}} \mathrm{d}x = 2\left(\int_t^\infty \mathrm{d}\mu\right)^{\frac{1}{2}} \leq 2\,\frac{\sqrt{B}}{\sqrt{G(t)}}, \tag{4.22}$$

by the definition of B. Combining (4.22) and (4.21) results in the desired upper bound.

For the lower bound assume that f is non-negative. We can bound the left hand side in (4.16) for fixed $r > 0$ from below by

$$\int_r^\infty \mathrm{d}\mu \left(\int_0^r f(t)\,\mathrm{d}t\right)^2 \leq \int_0^\infty \left(\int_0^x f(t)\,\mathrm{d}t\right)^2 \mu(\mathrm{d}x) \leq C\int_0^\infty |f(x)|^2\,\mu(\mathrm{d}x). \tag{4.23}$$

Now, we want to show that for any r holds

$$\int_r^\infty \mathrm{d}\mu \int_0^r \frac{1}{\mu(x)}\,\mathrm{d}x \leq C, \tag{4.24}$$

which implies $B \leq C$. In the case $\int_0^r \frac{1}{\mu(x)}\,\mathrm{d}x = 0$ the estimate (4.24) is immediate. In the case $\int_0^r \frac{1}{\mu(x)}\,\mathrm{d}x = \infty$ exists a function $f(x)$ with $\int_0^r |f(x)|^2\,\mathrm{d}\mu(x) < \infty$ and $\int_0^r f(x)\mathrm{d}x = \infty$. Then (4.23) implies that $\int_r^\infty \mathrm{d}\mu = 0$, which also yields (4.24). In the last case, namely $0 < \int_0^r \frac{1}{\mu(x)}\,\mathrm{d}x$, set $f(x) = \frac{1}{\mu(x)}$ and observe that (4.23) becomes

$$\int_r^\infty \mathrm{d}\mu \left(\int_0^r \frac{1}{\mu(x)}\,\mathrm{d}x\right)^2 \leq C\int_0^r \frac{1}{\mu(x)}\,\mathrm{d}x,$$

which gives, after dividing by $\int_0^r \frac{1}{\mu(x)}\mathrm{d}x$, the desired estimate (4.24). □

From the previous theorem it is easy to obtain the following proposition, which determines the Poncaré inequality in one dimension up to a factor of 8 for a general measure. The argument was found by [BG99], which considered the logarithmic Sobolev inequality in the continuous case and also by [Mic99], which considered the Poincaré inequality in the discrete case.

Proposition 4.16 (From Muckenhoupt functional to Poincaré inequality)**.** *Let μ be a probability measure on $\mathbb{R}$ absolute continuous w.r.t. to the Lebesgue measure and values in $\mathbb{R}^+ \cup \{+\infty\}$. Then the constants B_m^- and B_m^+ defined by*

$$B_m^- = \sup_{x\leq m}\left(\int_x^m \frac{1}{\mu(y)}\mathrm{d}y \int_{-\infty}^x \mathrm{d}\mu\right)$$
$$B_m^+ = \sup_{x\geq m}\left(\int_m^x \frac{1}{\mu(y)}\mathrm{d}y \int_x^\infty \mathrm{d}\mu\right)$$

are finite for μ-a.e. m if and only if they are finite for one common m.

Further, μ satisfies a Poincaré inequality PI(ϱ) if and only if B^-_m and B^+_m are finite. In this case ϱ obeys the estimate

$$\max\left\{(1-F_\mu(m))B^-_m, F_\mu(m)B^+_m\right\} \le \varrho^{-1} \le 4\max\left\{B^-_m, B^+_m\right\},$$

where $F_\mu(m) = \mu((-\infty, m])$.

Remark 4.17. A common cited version (see for instance [Fou05]) of the above proposition is given by setting $m = m^*$ with m^* equal to the median of μ, i.e. $\mu((-\infty, m^*]) = \frac{1}{2}$. In this case, the estimate of Proposition 4.16 has the form of

$$\frac{1}{2}\max\left\{B^-_{m^*}, B_{m^*}\right\} \le \varrho^{-1} \le 4\max\left\{B^-_{m^*}, B_{m^*}\right\}.$$

This gives a general characterization of the spectral gap or Poincaré constant up to a factor of 8. However, the median is hard to compute in general and the result stated in Proposition 4.16 is easier to obtain upper bounds.

Proof of Proposition 4.16. In the first step we will show that if m^* exists such that $B^-_{m^*}$ and $B^+_{m^*}$ are finite, then B^-_m and B^+_m are finite for μ-a.e. m. Therefore, we write $B^-_m(x) = \int_x^m \frac{1}{\mu(y)}\mathrm{d}y \int_{-\infty}^x \mathrm{d}\mu$ such that with this notation $B^-_m = \sup_{x\le m} B^-_m(x)$ and similarly for $B^+_m(x)$. By symmetry it is enough to consider $m \ge m^*$. Then, an immediate consequence is $B^+_m \le B^+_{m^*}$. To obtain a bound for $B^-_m(x)$ with $x \le m$, we split up the integral in the definition of $B^-_m(x)$ and obtain

$$\begin{aligned} B^-_m(x) &= \int_x^m \frac{1}{\mu(y)}\mathrm{d}y \int_{-\infty}^x \mathrm{d}\mu \le \mathbb{1}_{(-\infty,m^*]}(x)\int_x^{m^*} \frac{1}{\mu(y)}\mathrm{d}y\int_{-\infty}^x \mathrm{d}\mu + \int_{m^*}^m \frac{1}{\mu(y)}\mathrm{d}y\int_{-\infty}^x \mathrm{d}\mu \\ &\le \mathbb{1}_{(-\infty,m^*]}(x)B^-_{m^*}(x) + B^+_{m^*}(m)\frac{F_\mu(x)}{1-F_\mu(m)} \le B^-_{m^*} + B^+_{m^*}\frac{F_\mu(x)}{1-F_\mu(m)}, \end{aligned}$$

which is finite for m with $0 < F_\mu(m) < 1$, hence for μ-a.e. m.

Now, we want to deduce the upper bound for the inverse spectral gap, which follows by noting that for an arbitrary $m \in \mathbb{R}$ holds

$$\operatorname{var}_\mu(f) \le \int_{-\infty}^{\infty} (f - f(m))^2\,\mathrm{d}\mu = \int_{-\infty}^{m} (f-f(m))^2\,\mathrm{d}\mu^-_m + \int_m^\infty (f-f(m))^2\,\mathrm{d}\mu^+_m \tag{4.25}$$

where we introduced the measures

$$\mu^-_m(\mathrm{d}x) = \mathbb{1}_{(-\infty,m)}(x)\mu(\mathrm{d}x) \qquad \text{and} \qquad \mu^+_m = \mathbb{1}_{(m,+\infty)}(x)\mu(\mathrm{d}x).$$

We can apply Theorem 4.15 to both of the last integrals in (4.25) with the measure μ substituted by $\mu^\pm_m$ and obtain the estimate (4.16) with a constant $C^\pm_m$, which is bounded by $4B^\pm_m$. For μ^+_m this leads to

$$\int_m^\infty (f-f(m))^2\,\mathrm{d}\mu^+_m = \int_m^\infty \left(\int_m^y f'(x)\mathrm{d}x\right)^2 \mathrm{d}\mu^+_m(y) \le 4B^+_m \int_m^\infty \left|f'\right|^2 \mathrm{d}\mu^+_m,$$

and similarly for μ_m^-. This proves out right after again combining both integrals

$$\operatorname{var}_\mu(f) \leq 4B_m^- \int_{-\infty}^{m} \left|f'\right|^2 \mathrm{d}\mu_m^- + 4B_m^+ \int_{m}^{\infty} \left|f'\right|^2 \mathrm{d}\mu_m^+ \leq 4\max\left\{B_m^-, B_m^+\right\} \int_{-\infty}^{\infty} \left|f'\right|^2 \mathrm{d}\mu$$

the upper bound.

Once again we start with equation (4.25) and consider the integral over (m, ∞). Now we choose $0 < \tau < C_m^+$ and a function g_τ such that for the optimal constant C_m^+ in

$$\int_m^\infty (f - f(m))^2 \,\mathrm{d}\mu \leq C_m^+ \int_m^\infty \left|f'\right|^2 \mathrm{d}\mu$$

holds

$$\int_m^\infty (g_\tau - g_\tau(m))^2 \,\mathrm{d}\mu \geq \left(C_m^+ - \tau\right) \int_m^\infty \left|g_\tau'\right|^2 \mathrm{d}\mu. \tag{4.26}$$

These g_τ exists thanks to the optimality. By Theorem 4.15 we know that $B_m^+ \leq C_m^+$. We can assume w.l.o.g. $g_\tau(m) = 0$, else we could consider $\tilde{g}_\tau(x) = g_\tau(x) - g_\tau(m)$. Now, we set $g_\tau(x) = 0$ for $x < m$ and note $\mu(\{g_\tau = 0\}) \geq \mu((-\infty, m])) = F_\mu(m)$. Therewith, it follows

$$\left(\int g_\tau \mathrm{d}\mu\right)^2 \leq \mu(\{g_\tau \neq 0\}) \int g_\tau^2 \mathrm{d}\mu \leq (1 - F_\mu(m)) \int g_\tau^2 \mathrm{d}\mu.$$

This results in the estimate

$$\begin{aligned} \operatorname{var}_\mu(g_\tau) &= \int g_\tau^2 \mathrm{d}\mu - \left(\int g_\tau \mathrm{d}\mu\right)^2 \geq F_\mu(m) \int g_\tau^2 \mathrm{d}\mu \\ &\overset{(4.26)}{\geq} F_\mu(m)(C_m^+ - \tau) \int \left|g_\tau'\right|^2 \mathrm{d}\mu \geq F_\mu(m)(B_m^+ - \tau) \int \left|g_\tau'\right|^2 \mathrm{d}\mu. \end{aligned}$$

Hence, $\varrho^{-1} \geq F_\mu(m)(B_m^+ - \tau)$ and sending $\tau \to 0$ the desired result follows. The case for the integral on $(-\infty, m)$ follows similarly by symmetry. ☐

Now, we have completed all the preparatory work and can directly proceed to the proof Proposition 4.2, which is restated at this point.

Proposition 4.2. *Let $\mu(\cdot|\eta)$ be a radial-mixing measure in the sense of Definition 4.14. Then the measure $\mu(\cdot|\eta)$ satisfies $\mathrm{PI}(\varrho(\eta))$ with constant*

$$\frac{1}{\varrho(\eta)} = O(\varepsilon) \qquad \textit{for} \quad \varepsilon \to 0$$

uniformly in $\eta \in S^{n-1}$.

Proof of Proposition 4.2. We deduce the desired statement by an application of Proposition 4.16. Therefore, we have to show that $B_m^\pm = O(\varepsilon)$ for a particular choice of m.

For this purpose, let us set $m = \sqrt{\frac{\varepsilon}{h''_\eta(0)}}$. Additionally, we assume that ε is small enough such that $m \le \frac{\delta}{2}$ with $\frac{\delta}{2}$ given by the conditions ($\tilde{\mathbf{H}}\mathbf{1}$) and ($\tilde{\mathbf{H}}\mathbf{2}$).

The measure $\mu(\cdot|\eta)$ is absolutely continuous and has by denoting $h_\eta(r) := H(\psi_\eta(r)$ the form

$$\mu(\mathrm{d}r) = \frac{1}{\hat{\mu}(\eta)} j_\eta(r) e^{-\frac{h_\eta(r)}{\varepsilon}} \mathrm{d}r, \text{ where } \hat{\mu}(\eta) = \int j_\eta(r) e^{-\frac{h_\eta(r)}{\varepsilon}} \mathrm{d}r \text{ and } h_\eta(0) = h'_\eta(0) = 0.$$

Furthermore, by ($\tilde{\mathbf{H}}\mathbf{2}$) it holds $h''_\eta(0) = \langle \eta, \nabla^2 H(0)\eta \rangle \ge \lambda_{\min} > 0$ by the non-degenerate Assumption 1.3. We start with the estimation of $B^-_m(r)$ for $r \le m = O(\varepsilon^{\frac{1}{2}})$. Then, according to (**j1**) of Lemma 4.12 it holds $j_\eta(r) = r^{n-1}$. Now, we use the expansion $h_\eta(r) = h''_\eta(0)\frac{r^2}{2} + O(r^3) = h''_\eta(0)\frac{r^2}{2} + O(\varepsilon^{\frac{3}{2}})$ to find the estimate

$$\begin{aligned} B^-_m(r) &= \int_0^r j_\eta(t) \exp\left(-\frac{1}{\varepsilon} h_\eta(t)\right) \mathrm{d}t \int_r^m \frac{1}{j_\eta(t)} \exp\left(\frac{1}{\varepsilon} h_\eta(t)\right) \mathrm{d}t \\ &= \int_0^r t^{n-1} \exp\left(-\frac{h''_\eta(0)t^2}{2\varepsilon} + O(\sqrt{\varepsilon})\right) \mathrm{d}t \int_r^m t^{-n+1} \exp\left(\frac{h''_\eta(0)t^2}{2\varepsilon} + O(\sqrt{\varepsilon})\right) \mathrm{d}t \end{aligned}$$

We continue with the variable substitution $r = sm$ and $t = zm$ and arrive at

$$\begin{aligned} B^-_m(r) &\le m^2 \int_0^s z^{n-1} \exp\left(-\frac{z^2}{2}\right) \mathrm{d}z \int_s^1 z^{-n+1} \exp\left(\frac{z^2}{2}\right) \mathrm{d}z \left(1 + O(\sqrt{\varepsilon})\right) \\ &\le \varepsilon \frac{E_n}{h''_\eta(0)} (1 + O(\sqrt{\varepsilon})). \end{aligned}$$

The constant E_n is given as upper bound in the estimate

$$\int_0^s z^{n-1} \underbrace{e^{-\frac{z^2}{2}}}_{\le 1} \mathrm{d}z \int_s^1 z^{-n+1} \underbrace{e^{\frac{z^2}{2}}}_{\le \sqrt{e}} \mathrm{d}z \le \sqrt{e}\frac{s^n}{n} \begin{cases} \ln(\frac{1}{s}) & , n = 2 \\ \frac{1 - s^{n-2}}{(n-2)s^{n-2}} & , n > 2 \end{cases}.$$

Now, we bound the right hand side of the above estimate. In the case $n = 2$, the function $s \mapsto s^2 \ln \frac{1}{s}$ attains its maximum for $s = \frac{1}{\sqrt{e}}$ with value $\frac{1}{2e}$. Hence, we have $E_2 = \frac{1}{4\sqrt{e}}$. In the case $n > 2$, the function $s \mapsto s^2(1 - s^{n-2})$ attains its maximum for $s = \left(\frac{2}{n}\right)^{\frac{1}{n-2}}$ with value $\left(\frac{2}{n}\right)^{\frac{2}{n-2}} \left(1 - \frac{2}{n}\right) \le \frac{n-2}{n}$. Hence, we can set $E_n = \frac{\sqrt{e}}{n^2}$. We note that in any case holds $E_n \le 1$ independently of n.

Now, we continue with the estimation of $B^+_m(r)$ for $r \ge m$. Therefore, we can use the properties (**h1**) and (**h2**) of Proposition 4.13 as well as the fact the $j_\eta(\cdot)$ is non-zero in

$(0, T_\eta)$ by Lemma 4.12

$$\begin{aligned} B_m^+(r) &= \int_m^r \frac{1}{j_\eta(t)} e^{\frac{1}{\varepsilon} h_\eta(t)} \mathrm{d}t \int_r^{T_\eta} j_\eta(t) e^{-\frac{1}{\varepsilon} h_\eta(t)} \mathrm{d}t \\ &\leq \int_m^r \frac{1}{j_\eta(t)} e^{\frac{1}{\varepsilon} h_\eta(r) - \lambda_2 \frac{(r-t)^2}{\varepsilon}} \mathrm{d}t \int_r^{T_\eta} j_\eta(t) e^{-\frac{1}{\varepsilon} h_\eta(r) - \lambda_1 \frac{(r-t)^2}{\varepsilon}} \mathrm{d}t \\ &= \varepsilon \frac{\pi}{4\sqrt{\lambda_1 \lambda_2}} + O(\varepsilon^{\frac{3}{2}}), \end{aligned}$$

where we have used the Laplace method in the last step. □

4.4. A polar mean-difference estimate: proof of Proposition 4.3

This section is devoted to the proof of Proposition 4.3. We have to show the estimate

$$\iint \left(\hat{f}(\eta) - \hat{f}(\theta) \right)^2 \hat{\mu}(\mathrm{d}\eta) \hat{\mu}(\mathrm{d}\theta) \leq O(\varepsilon) \int |\nabla f|^2 \, \mathrm{d}\mu.$$

The procedure is similar to the proof of the mean-difference estimate of Theorem 1.9. Analogous to section 3.2, we show in the first step that it is sufficient to consider the mean-difference estimate w.r.t. the simpler measures $\nu(\cdot|\eta)$ that are Gaussian approximations of the measures $\mu(\cdot|\eta)$. This step is the content of Lemma 4.18. Following the ideas of Section 3.1, we estimate in the second step the mean-difference w.r.t. the approximations $\nu(\cdot|\eta)$ by using a transport argument. This step is the content of Lemma 4.19. Compared to Section 3.1, we have to argue more carefully since the support of the measures $\nu(\cdot|\eta)$ is only one-dimensional. Therefore, we need an additionally ingredient in the proof of Lemma 4.19: It is an identity for the spherical mean and is provided by Lemma 4.23.

Provided the ingredients are valid, the proof of Proposition 4.3 just consists of a straightforward application of Lemma 4.18 and Lemma 4.19.

Now, we turn to the first step. Let us introduce the approximation for the measures $\mu(\mathrm{d}r|\eta)$ and $\hat{\mu}(\mathrm{d}\eta)$. Recall that in Section 3.2 we introduced the truncated Gaussian measures ν_i on any domain of attraction Ω_i (cf. (3.6)). In this section we had the convention to omit the index i and considered Ω to be a domain of attraction. We also assumed w.l.o.g. that the unique local minima of H on Ω is located at $0 \in \Omega$. Using these conventions we may write the approximation ν of μ on Ω as

$$\nu(\mathrm{d}x) = \frac{1}{Z_\nu} \mathbb{1}_E(x) e^{-\frac{\nabla^2 H(0)[x]}{2\varepsilon}} \mathrm{d}x, \quad \text{with} \quad Z_\nu = \int_E e^{-\frac{\nabla^2 H(0)[x]}{2\varepsilon}} \mathrm{d}x,$$

where the set E is given by

$$E = \{x \in \mathbb{R}^n : |\sqrt{\nabla^2 H(0)} x| \leq \sqrt{2\varepsilon}\, \omega(\varepsilon)\}.$$

In the new coordinates given by $\psi_\eta(r)$, the measure $\nu(\mathrm{d}x)$ becomes (cf. (4.1))

$$\nu(\mathrm{d}r, \mathrm{d}\eta) = \frac{1}{Z_\nu}\, \mathbb{1}_E(\psi_\eta(r))\, r^{n-1}\, e^{-\frac{\nabla^2 H(0)[\eta] r^2}{2\varepsilon}}\, \mathrm{d}r\, \mathrm{d}\eta.$$

In the last identity we already applied the fact that for ε small enough the Jacobian determinant is given by $j_\eta(r) = r^{n-1}$ (cf. (**j1**) of Lemma 4.12). Following (4.2), (4.3) and (4.4), we decompose the measure $\nu(\mathrm{d}r, \mathrm{d}\eta)$ according to

$$\nu(\mathrm{d}r, \mathrm{d}\eta) = \nu(\mathrm{d}r|\eta)\, \hat{\nu}(\mathrm{d}\eta),$$

where the conditional measures $\nu(\mathrm{d}r|\eta)$ and the marginal $\hat{\nu}(\mathrm{d}\eta)$ are given by

$$\begin{aligned} \nu(\mathrm{d}r|\eta) &= \frac{1}{Z_\nu \hat{\nu}(\eta)}\, \mathbb{1}_E(\psi_\eta(r))\, r^{n-1}\, e^{-\frac{\nabla^2 H(0)[\eta] r^2}{2\varepsilon}}\, \mathrm{d}r \quad \text{and} \\ \hat{\nu}(\mathrm{d}\eta) &= \frac{1}{Z_\nu} \int_E r^{n-1}\, e^{-\frac{\nabla^2 H(0)[\eta] r^2}{2\varepsilon}}\, \mathrm{d}r\, \mathrm{d}\eta. \end{aligned}$$

Now, we are able to formulate the first ingredient of the proof of Proposition 4.3.

Lemma 4.18 (Approximation by truncated Gaussians.)**.** *Using the notation*

$$\hat{f}(\eta) := \int f(\psi_\eta(r))\mu(\mathrm{d}r|\eta) \qquad \textit{and} \qquad \tilde{f}(\eta) := \int f(\psi_\eta(r))\nu(\mathrm{d}r|\eta),$$

it holds for any function f that

$$\iint \left(\hat{f}(\eta) - \hat{f}(\theta)\right)^2 \hat{\mu}(\mathrm{d}\eta)\hat{\mu}(\mathrm{d}\theta) \leq 3 \iint \left(\tilde{f}(\eta) - \tilde{f}(\theta)\right)^2 \hat{\mu}(\mathrm{d}\eta)\hat{\mu}(\mathrm{d}\theta) + O(\varepsilon) \int |\nabla f|^2\, \mathrm{d}\mu.$$

Proof of Lemma 4.18. Using the triangle inequality we can estimate

$$\left(\hat{f}(\eta) - \hat{f}(\theta)\right)^2 \leq 3\left(\left(\hat{f}(\eta) - \tilde{f}(\eta)\right)^2 + \left(\tilde{f}(\eta) - \tilde{f}(\theta)\right)^2 + \left(\tilde{f}(\theta) - \hat{f}(\theta)\right)^2\right).$$

By applying the same strategy as in Lemma 3.6, we can estimate the first and third term of the last inequality by

$$\left(\hat{f}(\eta) - \tilde{f}(\eta)\right)^2 \leq \frac{\operatorname{var}_{\mu(\cdot|\eta)}\left(\frac{\mathrm{d}\nu(\cdot|\eta)}{\mathrm{d}\mu(\cdot|\eta)}\right)}{\varrho_\eta} \int |\partial_r f(\psi_\eta(r))|^2\, \mathrm{d}\mu(\mathrm{d}r|\eta),$$

where ϱ_η is the Poincaré constant of $\mu(\cdot|\eta)$. By Proposition 4.2, we know that

$$\varrho_\eta^{-1} = O(\varepsilon).$$

Additionally, one can show as in Lemma 3.8 that

$$\operatorname{var}_{\mu(\cdot|\eta)}\left(\frac{\mathrm{d}\nu(\cdot|\eta)}{\mathrm{d}\mu(\cdot|\eta)}\right) = O(\sqrt{\varepsilon}\, \omega^3(\varepsilon)).$$

Finally, we only have to express $\partial_r f$ in terms of the full gradient ∇f. Because of the parametrization of the coordinates $\psi_\eta(r)$ by arc-length, we have $|\dot{\psi}_\eta(r)| = 1$. This yields the identity

$$|\partial_r f(\psi_\eta(r))| \leq |\nabla f(\psi_\eta(r))| \, |\dot{\psi}_\eta(r)| = |\nabla f(\psi_\eta(r))| \, ,$$

which completes the argument. □

Now, let us turn to the second ingredient of the proof of Proposition 4.3, namely:

Lemma 4.19. *Using the notation*

$$\tilde{f}(\eta) := \int f(\psi_\eta(r)) \nu(\mathrm{d}r|\eta),$$

it holds for any function f that

$$\iint \left(\tilde{f}(\eta) - \tilde{f}(\theta) \right)^2 \mathrm{d}\hat{\mu}(\eta) \mathrm{d}\hat{\mu}(\theta) \leq O(\varepsilon) \int |\nabla f|^2 \, \mathrm{d}\mu. \tag{4.27}$$

The last statement is verified by using a *polar transport interpolation* argument. Before turning to the proof of Lemma 4.19, some preparatory work has to be done.

For the transport argument showing (4.27), we embed $\nu(\cdot|\eta)$ and $\mu(\cdot|\eta)$ along ηr as one-dimensional measures in $\mathbb{R}^n$ and note that $\nu(\cdot|\eta) \ll \mu(\cdot|\eta)$ for all $\eta \in S^{n-1}$. Then, we can still define for $\eta, \theta \in S^{n-1}$ a transport distance like $\mathcal{T}_\mu$ by considering the Radon-Nikodym derivative in the cost density w.r.t. $\mu(\cdot|\eta)$.

We denote by $\sphericalangle(\eta, \theta) \subset S^{n-1}$ the geodesic on S^{n-1} connecting η and θ parametrized on $[0, |\sphericalangle(\eta,\theta)|]$, where $|\sphericalangle(\eta,\theta)|$ is the arc-length of the geodesic bounded for any $\eta, \theta \in S^{n-1}$. Note that $\sphericalangle(\eta,\theta)$ is unique for almost all $\eta, \theta \in S^{n-1}$. Let $(\eta_s)_{s \in [0, |\sphericalangle(\eta,\theta)|]}$ be its constant speed parametrization. Then consider the transport interpolation defined on $\operatorname{supp} \nu(\cdot|\eta_0)$ by

$$\Phi_s(\eta_0 r) := \sigma_s^{\frac{1}{2}} \sigma_0^{-\frac{1}{2}} \eta_s r \qquad \text{where} \qquad \sigma_s^{-1} = \nabla^2 H(0)[\eta_s]. \tag{4.28}$$

The transport interpolation is chosen such that

$$(\Phi_s)_\sharp \nu(\cdot|\eta_0) = \nu(\cdot|\eta_s).$$

In the proof of Lemma 4.19 we need an estimate on the transport speed, namely:

Lemma 4.20. *With the definitions from above it holds*

$$|\dot{\Phi}_s \circ \Phi_s^{-1}(r\eta_s)| \leq r \sqrt{1 + \left(\frac{\lambda_{\max}}{\lambda_{\min}} \right)^2}, \tag{4.29}$$

where $\lambda_{\min}$ and $\lambda_{\max}$ is the largest and smallest eigenvalue of $\nabla^2 H(0)$, respectively.

Proof of Lemma 4.20. First, for the derivative and the inverse of Φ_s, we have that

$$\dot{\Phi}_s(\eta_0 r) = \dot{\eta}_s \sigma_s^{\frac{1}{2}} \sigma_0^{-\frac{1}{2}} r - \eta_s \frac{1}{2} \frac{\dot{\sigma}_s}{\sqrt{\sigma_s \sigma_0}} r \quad \text{and} \quad \Phi_s^{-1}(\eta_s r) = \eta_0 \sigma_s^{-\frac{1}{2}} \sigma_0^{\frac{1}{2}} r.$$

Hence, the composition is given by

$$\dot{\Phi}_s \circ \Phi_s^{-1}(\eta_s r) = \dot{\eta}_s r - \eta_s \frac{1}{2} \frac{\dot{\sigma}_s}{\sigma_s} r.$$

By using $\eta_s \perp \dot{\eta}_s$ and $|\eta_s| = |\dot{\eta}_s| = 1$ we obtain the estimate

$$\left| \dot{\Phi}_s \circ \Phi_s^{-1}(\eta_s r) \right| = r \sqrt{|\dot{\eta}_s|^2 + \frac{1}{4} |\eta_s|^2 \left| \frac{\dot{\sigma}_s}{\sigma_s} \right|^2} = r \sqrt{1 + \frac{1}{4} \left| \frac{\dot{\sigma}_s}{\sigma_s} \right|^2}.$$

The result follows from the definition of σ_s (4.28) and the estimate

$$\left| \frac{\dot{\sigma}_s}{\sigma_s} \right| = \frac{2 \left| \left\langle \eta_s, \nabla^2 H(0) \dot{\eta}_s \right\rangle \right|}{\nabla^2 H(0)[\eta_s]} \leq 2 \, \frac{\lambda_{\max}}{\lambda_{\min}}.$$

□

The next lemma contains the main contribution to the proof of Lemma 4.19. It is the estimation of the mean-difference $(\tilde{f}(\eta) - \tilde{f}(\theta))$ by applying a transport argument similar to the argument outlined in Section 3.1.

Lemma 4.21. *For any $\eta, \theta \in S^{n-1}$ it holds*

$$\left(\tilde{f}(\eta) - \tilde{f}(\theta) \right)^2 \leq O(\varepsilon) \int_{\sphericalangle(\eta,\theta)} \int |\nabla f(r\xi)| \; \mu(\mathrm{d}r|\xi) \; \mathcal{H}^1(\mathrm{d}\xi),$$

uniformly in η, θ. We want to recall that $\sphericalangle(\eta, \theta)$ denotes the geodesic on S^{n-1} connecting η and θ.

Proof of Lemma 4.21. We estimate the mean-difference in the same spirit as in (3.1), namely

$$\begin{aligned}\left(\tilde f(\eta)-\tilde f(\theta)\right)^2 &= \left(\int_0^{|\sphericalangle(\eta,\theta)|}\int \frac{\mathrm{d}f(\Phi_s(r\eta_0))}{\mathrm{d}s}\ \nu(\mathrm{d}r|\eta_0)\ \mathrm{d}s\right)^2\\ &\le \left(\int_0^{|\sphericalangle(\eta,\theta)|}\int |\nabla f(r\eta_s)|\,|\dot\Phi_s\circ\Phi_s^{-1}(r\eta_s)|\ \nu(\mathrm{d}r|\eta_s)\ \mathrm{d}s\right)^2\\ &\overset{(4.29)}{\le} \kappa_H\left(\int_0^{|\sphericalangle(\eta,\theta)|}\int |\nabla f(r\eta_s)|\ r\ \frac{\nu(\mathrm{d}r|\eta_s)}{\mu(\mathrm{d}r|\eta_s)}\ \mu(\mathrm{d}r|\eta_s)\ \mathrm{d}s\right)^2,\end{aligned}$$

where $\kappa_H = 1+\left(\frac{\lambda_{\max}}{\lambda_{\min}}\right)^2$ from (4.29). From this point, we estimate more along the lines of Remark 3.2 by using Cauchy-Schwarz w.r.t. $L^2(\mu(\mathrm{d}r|\eta_s))$ and Jensen's inequality for the outer integral

$$\begin{aligned}\left(\tilde f(\eta)-\tilde f(\theta)\right)^2 &\le \kappa_H\,|\sphericalangle(\eta,\theta)|\int_0^{|\sphericalangle(\eta,\theta)|}\int |\nabla f(r\eta_s)|^2\,\frac{\nu(\mathrm{d}r|\eta_s)}{\mu(\mathrm{d}r|\eta_s)}\ \mu(\mathrm{d}r|\eta_s)\int r^2\ \nu(\mathrm{d}r|\eta_s)\ \mathrm{d}s\\ &\approx \kappa_H\pi\int_0^{|\sphericalangle(\eta,\theta)|}\int |\nabla f(r\eta_s)|^2\,\mu(\mathrm{d}r|\eta_s)\int r^2\ \nu(\mathrm{d}r|\eta_s)\ \mathrm{d}s\end{aligned}$$

by using again the error estimate on the approximation given in Lemma 3.8 and therefore $\frac{\nu(\mathrm{d}r|\eta_s)}{\mu(\mathrm{d}r|\eta_s)} = 1+O(\sqrt{\varepsilon}|\ln^{\frac{3}{2}}(\varepsilon)|)$. Finally, note that the second moment of $\nu(r|\eta_s)$ is $O(\varepsilon)$ uniformly in η_s, due to the non-degeneracy assumption on the Hessian $\nabla^2 H(0)$ in the minimum (cf. Assumption 1.2). ☐

Before turning to the proof of Lemma 4.19 we need two more ingredients. The next lemma contains an asymptotic characterization of the relative density of the marginal measure $\hat\mu$ w.r.t. the uniform probability measure on S^{n-1} denoted by ς.

Lemma 4.22. *The relative density of $\hat\mu$ w.r.t. the uniform probability measure on S^{n-1} denoted by ς can be estimated for any $\xi\in S^{n-1}$ as*

$$\frac{\mathrm{d}\hat\mu}{\mathrm{d}\varsigma}(\xi)\approx\frac{\sqrt{\det\nabla^2 H(0)}}{(\nabla^2 H(0)[\xi])^{\frac{n}{2}}}.$$

In particular, the last identity yields for all $\xi\in S^{n-1}$ the uniform bound

$$\left(\frac{\lambda_{\max}}{\lambda_{\min}}\right)^{-\frac{n}{2}}\lesssim\frac{\mathrm{d}\hat\mu}{\mathrm{d}\varsigma}(\xi)\lesssim\left(\frac{\lambda_{\max}}{\lambda_{\min}}\right)^{\frac{n}{2}}. \tag{4.30}$$

Proof of Lemma 4.22. By the assumption (**$\tilde{\mathbf{H}}$2**) the coordinates are spherically around the local minimum. Hence, expanding H in 0 in the direction ξ yields

$$H(r\xi)=H(0)+\nabla^2H(0)[\xi]\frac{r^2}{2}+O(r^3).$$

Now, we choose Z_μ such that $H(0)=0$. Then it holds that

$$\begin{aligned}\hat{\mu}(\xi) &= \frac{1}{Z_\mu}\int_0^{T_\xi} e^{-\frac{H(\psi_\xi(r))}{\varepsilon}} j_\xi(r)\,\mathrm{d}r \\ &= \frac{1}{Z_\mu}\int_0^{\sqrt{\varepsilon}\,\omega(\varepsilon)} r^{n-1}\, e^{-\frac{\nabla^2 H(0)[\xi] r^2}{2\varepsilon}}\,\mathrm{d}r\,\big(1+O(\sqrt{\varepsilon}\,\omega^3(\varepsilon)\big) \;+\; \frac{1}{Z_\mu}\int_{\sqrt{\varepsilon}\,\omega(\varepsilon)}^{T_\xi} e^{-\frac{H(\psi_\xi(r))}{\varepsilon}}\,\mathrm{d}r \\ &= \frac{1}{Z_\mu}\frac{1}{2}\left(\frac{2\varepsilon}{\nabla^2 H(0)[\xi]}\right)^{\frac{n}{2}}\Gamma\left(\frac{n}{2}\right)\big(1+O(\sqrt{\varepsilon}\,\omega^3(\varepsilon)\big)+O(\exp^{-\omega^2(\varepsilon)}).\end{aligned}$$

By noting that $Z_\mu \approx (2\pi\varepsilon)^{\frac{n}{2}}\big(\det \nabla^2 H(0)\big)^{-\frac{1}{2}}$, we arrive at

$$\hat{\mu}(\xi) \approx \frac{\sqrt{\det \nabla^2 H(0)}}{(\nabla^2 H(0)[\xi])^{\frac{n}{2}}}\;\frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}} \approx \frac{\sqrt{\det \nabla^2 H(0)}}{(\nabla^2 H(0)[\xi])^{\frac{n}{2}}}\;\varsigma(\xi),$$

where we used the fact that the density $\varsigma(\xi)$ is given by the constant

$$\varsigma(\xi) = \big(\mathcal{H}^{n-1}(S^{n-1})\big)^{-1} = \frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}}.$$

□

The last remaining ingredient for the proof of Lemma 4.19 is the following basic statement.

Lemma 4.23 (Spherical mean)**.** *Let ς be the normalized uniform measure on S^{n-1}. Then it holds for all $F : S^{n-1} \to L^1(S^{n-1},\varsigma)$ that*

$$\int_{S^{n-1}}\int_{S^{n-1}}\int_{\sphericalangle(\theta,\eta)} F(\xi)\;\mathcal{H}^1(\mathrm{d}\xi)\;\varsigma(\mathrm{d}\eta)\;\varsigma(\mathrm{d}\theta) = \frac{\pi}{2}\int_{S^{n-1}} F(\theta)\;\varsigma(\mathrm{d}\theta).$$

Proof. Choose $e \in S^{n-1}$ fixed. Then by the transitivity of SO^n on S^{n-1}, exists for all $\theta \in S^{n-1}$ a rotation $g_\theta \in SO^n$ such that $g_\theta\theta = e$. Then, the result follows from the variable substitutions first $\xi \mapsto g_\theta\xi$ and second $\eta \mapsto g_\theta\eta$

$$\begin{aligned}\int\limits_{S^{n-1}}\int\limits_{S^{n-1}}\int\limits_{\sphericalangle(\theta,\eta)} F(\xi)\;\mathcal{H}^1(\mathrm{d}\xi)\;\varsigma(\mathrm{d}\eta)\;\varsigma(\mathrm{d}\theta) &= \int\limits_{S^{n-1}}\int\limits_{S^{n-1}}\int\limits_{\sphericalangle(e,g_\theta^{-1}\eta)} F(g_\theta\xi)\;\mathcal{H}^1(\mathrm{d}\xi)\;\varsigma(\mathrm{d}\eta)\;\varsigma(\mathrm{d}\theta) \\ &= \int\limits_{S^{n-1}}\int\limits_{S^{n-1}}\int\limits_{\sphericalangle(e,\eta)} F(g_\theta\xi)\;\mathcal{H}^1(\mathrm{d}\xi)\;\varsigma(\mathrm{d}\eta)\;\varsigma(\mathrm{d}\theta)\end{aligned}$$

By interchanging the order of integration from $\mathcal{H}^1(\mathrm{d}\xi)\;\varsigma(\mathrm{d}\eta)\;\varsigma(\mathrm{d}\theta)$ to $\varsigma(\mathrm{d}\theta)\;\mathcal{H}^1(\mathrm{d}\xi)\;\varsigma(\mathrm{d}\eta)$ we find with the transitivity of SO^n on S^{n-1} that

$$\int_{S^{n-1}} F(g_\theta\xi)\;\varsigma(\mathrm{d}\theta) = \int_{S^{n-1}} F(\theta)\;\varsigma(\mathrm{d}\theta), \quad \text{independently of } \xi.$$

The last step is to observe, that the remaining double integral is the mean length of a geodesic on S^{n-1}

$$\int_{S^{n-1}} \int_{\sphericalangle(e,\eta)} \mathcal{H}^1(\mathrm{d}\xi)\, \varsigma(\mathrm{d}\eta) = \int_{S^{n-1}} |\sphericalangle(e,\eta)|\, \varsigma(\mathrm{d}\eta) = \frac{\pi}{2}.$$

□

Finally, we can turn to the proof of Lemma 4.19

Proof of Lemma 4.19. Integrating the inequality of Lemma 4.21 w.r.t. $\hat{\mu}(\eta)$ and $\hat{\mu}(\theta)$ results in the estimate

$$\iint \left(\tilde{f}(\eta) - \tilde{f}(\theta)\right)^2 \hat{\mu}(\mathrm{d}\eta)\hat{\mu}(\mathrm{d}\theta) \leq O(\varepsilon) \iint \int_{\sphericalangle(\eta,\theta)} \int |\nabla f(r\xi)|^2 \,\mu(\mathrm{d}r|\xi)\; \mathcal{H}^1(\mathrm{d}\xi)\; \hat{\mu}(\mathrm{d}\eta)\hat{\mu}(\mathrm{d}\theta).$$

Hence, we arrive by setting $F(\xi) = \int |\nabla f(r\xi)|^2 \mu(\mathrm{d}r|\xi)$ at

$$\iint \left(\tilde{f}(\eta) - \tilde{f}(\theta)\right)^2 \hat{\mu}(\mathrm{d}\eta)\hat{\mu}(\mathrm{d}\theta) \leq O(\varepsilon) \iint \int_{\sphericalangle(\eta,\theta)} F(\xi)\; ?\mathcal{H}^1(\mathrm{d}\xi)\; \varsigma(\mathrm{d}\eta)\varsigma(\mathrm{d}\theta). \tag{4.31}$$

Applying Lemma 4.23 to (4.31) and changing the measure back with the help of (4.30) results into

$$\begin{aligned}\iint \left(\tilde{f}(\eta) - \tilde{f}(\theta)\right)^2 \hat{\mu}(\mathrm{d}\eta)\hat{\mu}(\mathrm{d}\theta) &\leq O(\varepsilon) \int \int |\nabla f(\psi_\eta(r))|^2\; \mu(\mathrm{d}r|\eta)\; \hat{\mu}(\mathrm{d}\eta)\\ &= O(\varepsilon) \int |\nabla f|^2 \,\mathrm{d}\mu,\end{aligned}$$

which yields the desired statement (4.27). □

A. Construction of the perturbed Hamiltonian $\tilde{H}$: proof of Proposition 4.1

Recalling the notation of Section 4, we only consider one basin of attraction Ω_i (cf. (1.6)) and therefore omit the index i. We assume w.l.o.g. that $0 \in \Omega$ is the local minimum of H in Ω. This part of the appendix is devoted to the proof of Proposition 4.1.

Proposition 4.1. *Assume that the basin of attraction Ω is bounded. Then, there exist coordinates $\{\psi_\eta\}_{\eta\in S^{n-1}}$ on Ω such that the conditional measures $\mu(\mathrm{d}r|\eta)$ are radial-mixing measures in the sense of Definition 4.14.*

Hence, we have to show that on a basin of attraction Ω (cf. (1.6)) exists a perturbation $\tilde{H}$ of the Hamiltonian H satisfying the conditions ($\tilde{\mathbf{H}}\mathbf{1}$)-($\tilde{\mathbf{H}}\mathbf{5}$):

(i) $\tilde{H}$ equals H up to small δ-neighborhoods around the critical points of H, i.e. by recalling that $\mathcal{S}$ is the set of critical points of H we have

$$\forall x \notin \bigcup_{y\in\mathcal{S}} B_\delta(y): \ \tilde{H}(x) = H(x). \tag{$\tilde{\mathbf{H}}\mathbf{1}$}$$

(ii) $\tilde{H}$ restricted to a ball around the local minima located in 0 of radius $\frac{\delta}{2} > 0$ is spherically symmetric and linear, i.e.

$$\forall x \in B_{\frac{\delta}{2}}(0): \ \tilde{H}(x) = \tilde{H}(0) + |x|\,. \tag{$\tilde{\mathbf{H}}\mathbf{2}$}$$

(iii) $|\nabla \tilde{H}|$ is uniformly bounded from below in $\Omega\backslash\{0\}$, i.e.

$$\exists c_{\tilde{H}} > 0, \forall x \in \Omega\backslash\{0\}: \ |\nabla \tilde{H}(x)| \geq c_{\tilde{H}}. \tag{$\tilde{\mathbf{H}}\mathbf{3}$}$$

(iv) Gradient flow coordinates $(\psi_\eta(r))_{\eta\in S^{n-1}}$ obtained from $\tilde{H}$ in the way described in Definition 4.9 satisfy for all $\eta \in S^{n-1}$

$$\forall r \in (\delta/2, T_\eta): \quad -C_{\tilde{H}} \max\left\{1, (T_\eta - r)^{-2}\right\} \leq \nabla\cdot\tilde{F}(\psi_\eta(r)) \leq C_{\tilde{H}}, \tag{$\tilde{\mathbf{H}}\mathbf{4}$}$$

(v) and

$$\forall r \in (0, T_\eta): \langle \nabla H(\psi_\eta(r)), \nabla\tilde{H}(\psi_\eta(r))\rangle \geq c_{\tilde{H}} \min\{1, r, T_\eta - r\} \,|\nabla\tilde{H}(\psi_\eta(r))|. \tag{$\tilde{\mathbf{H}}\mathbf{5}$}$$

Because of ($\tilde{\mathbf{H}}\mathbf{1}$), the perturbation $\tilde{H}$ is only allowed to differ from the Hamiltonian H on small neighborhoods around the critical points $z \in \Omega$ of H. Therefore, we can consider every critical point separately and do an explicit local construction of the perturbation $\tilde{H}$ on these neighborhoods of critical points. In Section A.1, we consider the case where the critical point is a local extrema of H. In Section A.2, we consider the case that the critical point is a saddle point of H.

A.1. Correction at local extrema

In this section, we can omit the case where the critical point z is a local maxima. The reason is that the construction of $\tilde{H}$ would be very similar to the construction of $\tilde{H}$ around a local minima by considering $-H$ instead of H.

We denote by $\lambda_{\min}$ the smallest eigenvalue of $\nabla^2 H(0)$, which is strictly positive by the non-degeneracy Assumption 1.2. Hence, we can choose $\delta > 0$ such that H is strictly convex on $B_\delta(0)$. Let $\xi \in C^\infty(\mathbb{R}^+, [0,1])$ be a decreasing smooth step function satisfying the conditions

$$\forall r \in [0, 1/2]: \xi(r) = 1, \quad \forall r \in (1/2, 1): \xi'(r) < 0, \quad \text{and} \quad \forall r \in [1, \infty): \xi(r) = 0. \tag{A.1}$$

For convenience, we set $\xi_\delta(r) = \xi(\frac{r}{\delta})$. We define the correction $\tilde{H}$ of H around the minimum as

$$\tilde{H}(x) := \xi_\delta(|x|)\ (|x| - \delta) + (1 - \xi_\delta(|x|))\ H(x), \qquad \text{for } x \in B_\delta(0). \tag{A.2}$$



Figure 3: Contours and flow lines derived from H (left) and $\tilde{H}$ (right).

Now, we state the main result of this section.

Lemma A.1. *For δ small enough, the function $\tilde{H}$ satisfies the properties ($\tilde{\mathbf{H}}\mathbf{1}$)-($\tilde{\mathbf{H}}\mathbf{5}$) in the neighborhood $B_\delta(0)$ around the minimum $0 \in \Omega$.*

Proof. The property ($\tilde{\mathbf{H}}\mathbf{1}$) is fulfilled by construction.
The property ($\tilde{\mathbf{H}}\mathbf{2}$) is satisfied by the definition (A.2) of $\tilde{H}$ and the definition (A.1) of ξ, which ensures that the gradient-flow coordinates of $\tilde{H}$ coincide with polar coordinates on the ball $B_{\delta/2}(0)$.
The property ($\tilde{\mathbf{H}}\mathbf{3}$) is verified by a straight forward argument, which we outline in full details for the convenience of the reader. Direct calculation leads to the identity

$$\begin{aligned} &\nabla\tilde{H}(x) = a(x)\ \frac{x}{|x|} + b(x)\ \nabla H(x), \quad \text{with} \\ &a(x) = \delta^{-1}\xi_\delta'(|x|)\ (|x| - \delta - H(x)) + \xi_\delta(|x|) \quad \text{and} \quad b(x) = 1 - \xi_\delta(|x|). \end{aligned} \tag{A.3}$$

For the verification of ($\tilde{\mathbf{H}}\mathbf{3}$), we will need two observations.
The first one is $a(x) > \xi_\delta(|x|)$, which follows for small enough δ from our assumptions on H, namely $H(0) = 0$ and $H(x) \geq 0$ as well as the monotonicity (A.1) of ξ, i.e. $\xi' \leq 0$.
The second observation is that for $x \in B_\delta(0)$ it holds

$$\left\langle \frac{x}{|x|}, \nabla H(x) \right\rangle = |x|\ \left\langle \frac{x}{|x|}, \nabla^2 H(0)\ \frac{x}{|x|} \right\rangle + O(x^2) \geq \lambda_{\min}|x| + O(\delta^2) \geq \frac{\lambda_{\min}}{2}\,|x|\,, \tag{A.4}$$

which follows directly from Taylor expansion and choosing δ sufficiently small. In the last inequality, $\lambda_{\min} > 0$ denotes the smallest eigenvalue of $\nabla^2 H(0)$.
Let us now verify (**$\tilde{\mathbf{H}}$3**). The identity (A.3) directly yields for $x \in B_\delta(0)$

$$\begin{aligned}|\nabla \tilde{H}(x)| &= a^2(x) + 2\ a(x)\ b(x)\ \left\langle \frac{x}{|x|}, \nabla^2 H(x) \right\rangle + b^2(x)\ |\nabla H(x)|^2 \\ &\overset{\text{(A.4)}}{\geq} a^2(x) + 2\ a(x)\ b(x)\ \frac{\lambda_{\min}}{2} |x| + b^2(x)\ |\nabla H(x)|^2.\end{aligned}$$

We consider two cases. In the first case, we assume that $|x| < \frac{\delta}{4}$. Then we get

$$|\nabla \tilde{H}(x)| \geq a^2(x) + O(\delta^2) \overset{\text{(A.1)}}{\geq} \xi\,(1/4) + O(\delta^2) \geq c_{\tilde{H}} > 0.$$

In the second case, namely $\frac{\delta}{4} \leq |x| \leq \delta$, we get

$$\begin{aligned}|\nabla \tilde{H}(x)| &\geq b^2(x)\ |\nabla H(x)|^2 + O(\delta^2) \\ &\overset{\text{(A.1)}}{\geq} \left(1 - \xi\left(\tfrac{1}{4}\right)\right)^2 \min_{\frac{\delta}{4} \leq |x| \leq \delta} |\nabla H(x)|^2 + O(\delta^2) \geq c_{\tilde{H}} > 0,\end{aligned}$$

which yields the desired property (**$\tilde{\mathbf{H}}$3**).

Now, let us turn to the property (**$\tilde{\mathbf{H}}$4**). It suffices to show that for $x \in B_\delta(0) \backslash B_{\delta/2}(0)$ holds

$$\left| \nabla \cdot \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|}(x) \right| \leq C.$$

However, this easily follows from the observation that

$$\begin{aligned}\left| \nabla \cdot \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|}(x) \right| &= \frac{1}{|\nabla \tilde{H}(x)|}\ \left| \Delta \tilde{H}(x) - \left\langle \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|}(x), \nabla^2 \tilde{H}(x)\ \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|}(x) \right\rangle \right| \\ &\overset{(\tilde{\mathbf{H}}3)}{\leq} \frac{n-1}{c}\ \|\nabla^2 \tilde{H}(x)\|\end{aligned}$$

and the fact that $\tilde{H}(x)$ is by construction a smooth function on $B_\delta(0) \backslash B_{\delta/2}(0)$.

Finally, let us verify the property (**$\tilde{\mathbf{H}}$5**). It is sufficient to show that

$$\forall t \text{ such that } \psi_\eta(t) \in B_\delta(0): \quad \left\langle \nabla H(\psi_\eta(t)), \nabla \tilde{H}(\psi_\eta(t)) \right\rangle \geq c_{\tilde{H}}\ t\ |\nabla \tilde{H}(\psi_\eta(t))|,$$

where the local coordinates $\psi_\eta(t)$ are given by (4.14). Using (A.3) we get

$$\begin{aligned}\left\langle \nabla H(x), \nabla \tilde{H}(x) \right\rangle &= a(x)\ \left\langle \nabla H(x), \frac{x}{|x|} \right\rangle + b(x)\ |\nabla H(x)|^2 \\ &\geq a(x)\ \left\langle \nabla H(x), \frac{x}{|x|} \right\rangle + b(x)\ |\nabla H(x)|\ \left\langle \nabla H(x), \frac{x}{|x|} \right\rangle \\ &\overset{\text{(A.4)}}{\geq} (a(x) + b(x)\ |\nabla H(x)|)\ \frac{\lambda_{\min}}{2} |x| \\ &\overset{\text{(A.3)}}{\geq} \frac{\lambda_{\min}}{2}\ |x|\ |\nabla \tilde{H}(x)|.\end{aligned}$$

By the last inequality, it is only left to show that for some constant $c > 0$

$$|\psi_\eta(t)| \geq c\, t, \qquad \text{for all } \psi_\eta(t) \in B_\delta(0). \tag{A.5}$$

It follows from the definition (4.14) of $\psi_\eta(t)$ that

$$|\psi_\eta(t)| = \int_0^t \partial_s |\psi_\eta(s)|\, ds = \int_0^t \left\langle \frac{\psi_\eta(s)}{|\psi_\eta(s)|}, \frac{\nabla \tilde{H}(\psi_\eta(s))}{|\nabla \tilde{H}(\psi_\eta(s))|} \right\rangle ds. \tag{A.6}$$

For $\psi_\eta(s) \in B_\delta(0)$ it follows from (A.3) that $|\nabla \tilde{H}(\psi_\eta(s))| \leq C$. Therefore, we can estimate

$$\begin{aligned} &\left\langle \frac{\psi_\eta(s)}{|\psi_\eta(s)|}, \frac{\nabla \tilde{H}(\psi_\eta(s))}{|\nabla \tilde{H}(\psi_\eta(s))|} \right\rangle \\ &\quad \overset{\text{(A.3)}}{=} \frac{1}{|\nabla \tilde{H}(\psi_\eta(s))|} \left(a(\psi_\eta(s)) + b(\psi_\eta(s)) \left\langle \frac{\psi_\eta(s)}{|\psi_\eta(s)|}, \nabla H(\psi_\eta(s)) \right\rangle \right) \\ &\quad \overset{\text{(A.4)}}{\geq} \frac{1}{C} \left(a(\psi_\eta(s)) + b(\psi_\eta(s)) \frac{\lambda_{\min}}{2} |\psi_\eta(s)| \right) \\ &\quad \geq c_{\tilde{H}} > 0, \end{aligned} \tag{A.7}$$

where the last line is deduced with the same argument as used for (**$\tilde{\mathbf{H}}$3**). A combination of (A.6) and (A.7) yields the desired estimate (A.5) and therefore concludes the proof. □

A.2. Correction at the saddles

In this section, we give a construction of a perturbation $\tilde{H}$ satisfying Proposition 4.1 nearby saddle points. Every k-saddle p lies on the boundary of Ω. We will locally deform H in such a way that either trajectories will leave Ω nearby the saddle or they will avoid coming close to the saddle (cf. Figure 4). We achive this behavior by introducing a kink on the stable manifold (cf. Figure 5).

We assume w.l.o.g. that the saddle point p lies at the origin 0 and let Q be an orthogonal matrix, such that

$$\nabla^2 H(0) = Q \operatorname{diag}(\lambda_1^-, \ldots, \lambda_k^-, \lambda_1^+, \ldots, \lambda_{n-k}^+) Q^\top,$$

where $\lambda_1^-, \ldots, \lambda_k^- < 0$ and $\lambda_1^+, \ldots, \lambda_{n-k}^+ > 0$. Therefore, the stable $\mathcal{E}^s$ and unstable $\mathcal{E}^u$ subspaces of $\nabla^2 H(0)$ are given by

$$\begin{aligned} \mathcal{E}^{\mathrm{u}} &:= \left\{ Q(y,0)^\top \in \mathbb{R}^n : y \in \mathbb{R}^k \right\} \\ \mathcal{E}^{\mathrm{s}} &:= \left\{ Q(0,y)^\top \in \mathbb{R}^n : y \in \mathbb{R}^{n-k} \right\}. \end{aligned}$$


W^{u}
$\mathcal{E}_{\mathrm{u}}$
W^{s}
x_{s}
$\mathcal{E}_{\mathrm{s}}$
$h_{\mathrm{s}}(x_{\mathrm{s}})$
x_{u}
x



Figure 4: Flow lines derived from H (left) and $\tilde{H}$ (right).

The stable and unstable manifold of the saddle at 0 are given by

$$\begin{aligned} W^{\mathrm{s}} &:= \{y_0 \in \mathbb{R}^n : \dot{y}_t = -\nabla H(y_t), y_t \to 0\} \\ W^{\mathrm{u}} &:= \{y_0 \in \mathbb{R}^n : \dot{y}_t = \nabla H(y_t), y_t \to 0\}\,. \end{aligned}$$

Theorem A.2 (Stable manifold theorem [Tes12, Theorem 9.4, p. 259])**.** *For a small neighborhood $\mathcal{U}$ of the saddle 0 the locally stable manifold $W^{\mathrm{s}}_{\mathrm{loc}} := W^{\mathrm{s}} \cap \mathcal{U}$ are smooth manifolds tangent to $\mathcal{E}^{\mathrm{s}}$ at the origin. There exists neighborhoods $\mathcal{U}^{\mathrm{s}} \subseteq \mathcal{E}^{\mathrm{s}}$, $\mathcal{U}^{\mathrm{u}} \subseteq \mathcal{E}^{\mathrm{u}}$ and a smooth map $h_{\mathtt{s}} : \mathcal{U}^{\mathrm{s}} \to \mathcal{U}^{\mathrm{u}}$ satisfying*

$$h_{\mathtt{s}}(0) = 0 \quad \textit{and} \quad Dh_{\mathtt{s}}(0) = 0 \tag{A.8}$$

such that the local stable manifold allows the representation

$$W^{\mathrm{s}}_{\mathrm{loc}} = \{x_{\mathtt{s}} + h_{\mathtt{s}}(x_{\mathtt{s}}) : x_{\mathtt{s}} \in \mathcal{U}^{\mathrm{s}}\}\,.$$

Therewith, *local coordinates* on the saddle are given by the mapping

$$\xi(x_{\mathtt{s}}, x_{\mathtt{u}}) : \mathcal{U}^{\mathrm{s}} \times \mathcal{U}^{\mathrm{u}} \to \mathcal{U} \subset \mathbb{R}^n \qquad \xi(x_{\mathtt{s}}, x_{\mathtt{u}}) \mapsto x_{\mathtt{s}} + h_s(x_{\mathtt{s}}) + x_{\mathtt{u}}.$$

With the projections $\Pi_{\mathtt{s}}$ and $\Pi_{\mathtt{u}}$

$$\begin{aligned} \Pi_{\mathtt{s}} : \mathcal{U} \to \mathcal{U}^{\mathrm{s}} \qquad & \Pi_{\mathtt{s}} = Q\,\mathrm{diag}(\underbrace{0,\dots,0}_{k\text{-times}},\ \underbrace{1\dots,1}_{(n-k)\text{-times}}\ )Q^\top \\ \Pi_{\mathtt{u}} : \mathcal{U} \to \mathcal{U}^{\mathrm{u}} \qquad & \Pi_{\mathtt{u}} = Q\,\mathrm{diag}(\underbrace{1,\dots,1}_{k\text{-times}},\ \underbrace{0\dots,0}_{(n-k)\text{-times}}\ )Q^\top \end{aligned}$$

we obtain the reverse coordinate transformation by

$$\mathcal{U} \ni x \mapsto (\Pi_{\mathtt{s}} x, \Pi_{\mathtt{u}} x - h_{\mathtt{s}}(\Pi_{\mathtt{s}} x)) \in \mathcal{U}^{\mathtt{s}} \times \mathcal{U}^{\mathtt{u}}.$$

The construction of an additive perturbation is done with the help of two functions $p, q \in C^3(\mathbb{R}^+, [0,1])$ with support in $[0,1]$ such that

$$p(0) = 1, \quad p'(0) = 0, \quad \forall r \in (0,1) : p'(r) < 0 \tag{A$_p$}$$

$$\left|p'(r)\right| \leq 8r \text{ for } r \in [0,1] \tag{B$_p$}$$

$$p(r) \geq (1-2r)_+ \text{ for } r \in [0,1] \tag{C$_p$}$$

$$0 \leq r < s \leq 1 : \frac{p(r)}{\int_r^s p(t)\mathrm{d}t} \leq \frac{C_p}{(s-r)^2} \tag{D$_p$}$$

$$q(0) = 1, \quad \forall r \in (0,1) : q'(r) < 0 \tag{A$_q$}$$

$$(1-2r)_+ \leq \left|q'(r)\right| \leq 2 \text{ for } r \in [0,1] \tag{B$_q$}$$

Example A.3. The following both functions satisfy (A$_p$)-(D$_p$) and (A$_q$)-(B$_q$)

$$p(r) = (1-r^2)^3 \mathbb{1}_{[0,1)}(r) \qquad \text{and} \qquad q(r) = (1-r)(1-r^2)^2 \mathbb{1}_{[0,1)}(r).$$

Lemma A.4. *For constants $a, \delta > 0$ define the additive perturbation*

$$H^+_{\mathrm{loc}} : \mathbb{R}^+ \times \mathbb{R}^+ \to \mathbb{R} \qquad H^+_{\mathrm{loc}}(s,u) = a\delta^2 p\left(\frac{s}{\delta}\right) q\left(\frac{u}{\delta}\right), \tag{A.9}$$

where p and q satisfy (A$_p$)-(D$_p$) and (A$_q$)-(B$_q$), respectively. Let us define the modified Hamiltonian $\tilde{H}$ on $\mathcal{U} := \{x \in \mathbb{R}^n : |\Pi_s x| < \delta, |\Pi_u x - h_s(\Pi_s x)| < \delta\}$ by

$$\tilde{H}(x) := H(x) + H^+_{\mathrm{loc}}(|\Pi_s x|, |\Pi_u x - h_s(\Pi_s x)|). \tag{A.10}$$

*Then, there exists a constant $a > 0$ and δ small enough, such that $\tilde{H}$ on $\mathcal{U}$ satisfies the properties (**$\tilde{\mathbf{H}}$1**)-(**$\tilde{\mathbf{H}}$5**) of Assumption 4.8 and Theorem 4.1.*

Proof. The properties (**$\tilde{\mathbf{H}}$1**) and (**$\tilde{\mathbf{H}}$2**) are fulfilled by the construction of H^+_{loc} in (A.9).

Argument for (**$\tilde{\mathbf{H}}$3**): Let $x = x_{\mathtt{s}} + x_{\mathtt{u}} \in \mathcal{U} \backslash W^{\mathtt{s}}_{\mathrm{loc}}$ with $x_{\mathtt{s}} = \Pi_{\mathtt{s}} x$ and $x_{\mathtt{u}} = \Pi_{\mathtt{u}}$, hence $x_{\mathtt{u}} \neq h_{\mathtt{s}}(x_{\mathtt{s}})$. Further, set $s = |x_{\mathtt{s}}|$ and $u = |x_{\mathtt{u}} - h(x_{\mathtt{s}})|$. Then, it follows from (A.10)

$$\nabla \tilde{H}(x) = \nabla H(x) + \partial_s H^+_{\mathrm{loc}}(s,u) \frac{x_{\mathtt{s}}}{s} + \partial_u H^+_{\mathrm{loc}}(s,u)(\mathrm{Id} - Dh^\top_{\mathtt{s}}(x_{\mathtt{s}})) \frac{x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})}{u}.$$

Firstly, we expand $\nabla H(x)$ around the saddle point and find

$$\nabla H(x) = \nabla H(x_s + x_u) = \nabla^2 H(0) x_s + \nabla^2 H(0)(x_u - h_s(x_s)) + R_2(x), \tag{A.11}$$



Figure 5: $\tilde{H}$ (left) with a kink on the stable manifold and H^{+}_{loc} (right).

where the remainder term $R_2(x)$ satisfies $|R_2(x)| \leq (C_H + C_h)\,|x|^2 \leq (C_H + C_h)\delta^2$ as $|h_{\mathtt{s}}(x_{\mathtt{s}})| \leq C_h\,|x_{\mathtt{s}}|^2$ by (A.8). We note that the partial derivatives of the additive perturbation are given by

$$\partial_s H^{+}_{\mathrm{loc}}(s,u) = a\delta p'\left(\frac{s}{\delta}\right) q\left(\frac{u}{\delta}\right) \quad \text{and} \quad \partial_u H(s,u) = a\delta p\left(\frac{s}{\delta}\right) q'\left(\frac{u}{\delta}\right).$$

Combining all expressions for the gradient of the perturbed Hamiltonian results in

$$\begin{aligned}\nabla\tilde{H}(x) &= \left(\nabla^2 H(0)s + a\delta p'\left(\frac{s}{\delta}\right) q\left(\frac{u}{\delta}\right)\mathrm{Id}\right)\frac{x_{\mathtt{s}}}{s} + a\delta p\left(\frac{s}{\delta}\right) q'\left(\frac{u}{\delta}\right) Dh_{\mathtt{s}}^{\top}(x_{\mathtt{s}})\frac{x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})}{u} \\ &\qquad + \left(\nabla^2 H(0)u + a\delta p\left(\frac{s}{\delta}\right) q'\left(\frac{u}{\delta}\right)\mathrm{Id}\right)\frac{x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})}{u} + R_2(x) \\ &\quad =: N_{\mathtt{s}}(x) + N_{\mathtt{u}}(x) + R_2(x).\end{aligned} \tag{A.12}$$

We want to stress out that $\langle N_{\mathtt{s}}(x), N_{\mathtt{u}}(x)\rangle = 0$, because of $\nabla^2 H(0)$ leaves $\mathcal{E}^s$ and $\mathcal{E}^u$ invariant as well as $Dh_{\mathtt{s}}^{\top}(x_{\mathtt{s}})(x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})) \in \mathcal{U}^{\mathtt{s}}$ and, of course, also $\langle x_{\mathtt{s}}, x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})\rangle = 0$. This leads us to the conclusion

$$|\nabla\tilde{H}(x)|^2 = |N_{\mathtt{s}}(x)|^2 + |N_{\mathtt{u}}(x)|^2 + O(\delta^3) \geq \max\left\{|N_{\mathtt{s}}(x)|^2, |N_{\mathtt{u}}(x)|^2\right\} + O(\delta^3). \tag{A.13}$$

We proceed with the calculation of lower bounds for $|N_{\mathtt{s}}(x)|$ and $|N_{\mathtt{u}}(x)|$. We start with $|N_{\mathtt{s}}(x)|$ and denote with $\lambda^{+}_{\mathrm{min}}$ of the smallest positive eigenvalue of the Hessian $\nabla^2 H(0)$

$$|N_{\mathtt{s}}(x)| \geq \lambda^{+}_{\mathrm{min}} s - a\delta\left|p'\left(\frac{s}{\delta}\right)\right| q\left(\frac{u}{\delta}\right) - a\delta p\left(\frac{s}{\delta}\right) q'\left(\frac{u}{\delta}\right)|Dh_{\mathtt{s}}^{\top}(x_{\mathtt{s}})|. \tag{A.14}$$

We employ now the assumptions (B_p) and (A_q) to bound the second term in (A.14) by

$$a\delta\left|p'\left(\frac{s}{\delta}\right)\right| q\left(\frac{u}{\delta}\right) \leq 8as. \tag{A.15}$$

Likewise, we use the assumptions (A_p) and (B_q) and the additional observation that $\left|Dh_{\mathtt{s}}^{\top}(x_{\mathtt{s}})\right| \leq C_h\,|x_{\mathtt{s}}|$ by (A.8) to bound the third term in (A.14) by

$$a\delta p\left(\frac{s}{\delta}\right) q'\left(\frac{u}{\delta}\right)|Dh_{\mathtt{s}}^{\top}(x_{\mathtt{s}})| \leq 2a\delta C_h s \leq 2as, \qquad \text{for } \delta \leq C_h^{-1}. \tag{A.16}$$

Hence, we can deduce a lower bound for $|N_{\mathtt{s}}(x)|$ by choosing a small enough. Indeed, we observe

$$|N_{\mathtt{s}}(x)| \geq \left(\lambda_{\min}^{+} - 8a - 2a\right) s \geq \delta \, \frac{\lambda_{\min}^{+}}{2} \frac{s}{\delta} \quad \text{for} \quad a \leq \frac{\lambda_{\min}^{+}}{20}. \tag{A.17}$$

For the estimation of $|N_{\mathtt{u}}(x)|$, we use the fact that by the definition of $\mathcal{U}^{\mathtt{u}}$ the Hessian $\nabla^2 H(0)$ is strictly negative definite on $\mathcal{U}^{\mathtt{u}}$. Therefore, let us denote by $\lambda_{\min}^{-}$ the smallest negative eigenvalue of $\nabla^2 H(0)$ in modulus. In addition, we use the assumptions (C_p) for p and (B_q) for q to deduce the lower bound for $|N_{\mathtt{u}}(x)|$

$$\begin{aligned} |N_{\mathtt{u}}(x)| &\geq |\lambda_{\min}^{-}| u + a\delta p\left(\frac{s}{\delta}\right) \left| q'\left(\frac{s}{\delta}\right)\right| \overset{(\mathrm{C}_p),(\mathrm{B}_q)}{\geq} |\lambda_{\min}^{-}| u + a\delta \left(1 - 2\frac{s}{\delta}\right)_{+} \left(1 - 2\frac{u}{\delta}\right)_{+} \\ &\geq \min\left\{\frac{|\lambda_{\min}^{-}|\delta}{2}, a\delta\left(1 - 2\frac{s}{\delta}\right)_{+}\right\}, \end{aligned} \tag{A.18}$$

where the last step follows from either setting $u = 0$ or $u = \frac{\delta}{2}$. We combine the estimates (A.17) and (A.18) with (A.13) and arrive at

$$\begin{aligned} |\nabla \tilde{H}(x)| &\geq \delta \max\left\{\frac{\lambda_{\min}^{+}}{2}\frac{s}{\delta}, \min\left\{\frac{|\lambda_{\min}^{-}|}{2}, a\left(1 - \frac{2s}{\delta}\right)_{+}\right\}\right\} + O(\delta^{\frac{3}{2}}) \\ &= \delta \min\left\{\max\left\{\frac{\lambda_{\min}^{+}}{2}\frac{s}{\delta}, \frac{|\lambda_{\min}^{-}|}{2}\right\}, \max\left\{\frac{\lambda_{\min}^{+}}{2}\frac{s}{\delta}, a\left(1 - \frac{2s}{\delta}\right)_{+}\right\}\right\} + O(\delta^{\frac{3}{2}}) \\ &\geq \frac{\delta}{2} \min\left\{|\lambda_{\min}^{-}|, \max\left\{\lambda_{\min}^{+}\frac{s}{\delta}, a\left(1 - \frac{2s}{\delta}\right)_{+}\right\}\right\} + O(\delta^{\frac{3}{2}}) \\ &\geq \frac{\delta}{2} \min\left\{|\lambda_{\min}^{-}|, \frac{a}{\lambda_{\min}^{+} + 2a}\right\} + O(\delta^{\frac{3}{2}}) \end{aligned}$$

It holds for $a \leq \frac{\lambda_{\min}^{+}}{20}$ that $\frac{a}{\lambda_{\min}^{+} + 2a} \geq \frac{a}{2\lambda_{\min}^{+}}$. Therewith, we can assume w.l.o.g. that $a \leq \min\left\{\frac{\lambda_{\min}^{+}}{20}, 2|\lambda_{\min}^{-}|\lambda_{\min}^{+}\right\}$ and have the estimate

$$|\nabla \tilde{H}(x)| \geq \frac{\delta}{2}\frac{a}{2\lambda_{\min}^{+}} + O(\delta^{\frac{3}{2}}).$$

Hence, in the last step we can choose some δ small enough such that the error term $O(\delta^{\frac{3}{2}})$ can be absorbed

$$|\nabla \tilde{H}(x)| \geq \frac{\delta}{4}\frac{a}{2\lambda_{\min}^{+}} \quad \text{for} \quad a \leq \min\left\{\frac{\lambda_{\min}^{+}}{20}, 2\lambda_{\min}^{-}\lambda_{\min}^{+}\right\}.$$

Argument for ($\tilde{\mathbf{H}}\mathbf{4}$): Let us recall of what condition ($\tilde{\mathbf{H}}\mathbf{4}$) consists of. We have to deduce a lower and upper bound for

$$\nabla \cdot \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|} = \frac{1}{|\nabla \tilde{H}|}\left(\Delta \tilde{H} - \left\langle \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|}, \nabla^2 \tilde{H} \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|}\right\rangle\right).$$

We observe that if $\tilde{\lambda}_1(x) \leq \ldots \tilde{\lambda}_n(x)$ denote the ordered eigenvalues of $\nabla^2 \tilde{H}(x)$, then the following bound holds

$$(n-1)\tilde{\lambda}_1(x) \leq \sum_{i=1}^{n-1} \tilde{\lambda}_i(x) \leq \Delta \tilde{H} - \left\langle \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|}, \nabla^2 \tilde{H} \frac{\nabla \tilde{H}}{|\nabla \tilde{H}|} \right\rangle \leq \sum_{i=2}^{n} \tilde{\lambda}_i(x) \leq (n-1)\tilde{\lambda}_n(x). \tag{A.19}$$

For the estimation of $\tilde{\lambda}_1(x)$ and $\tilde{\lambda}_n(x)$, we use the variational formulation of the eigenvalues

$$\tilde{\lambda}_1(x) = \inf_{\eta \in S^{n-1}} \left\langle \eta, \nabla^2 \tilde{H}(x) \eta \right\rangle \qquad \text{and} \qquad \tilde{\lambda}_n(x) = \sup_{\eta \in S^{n-1}} \left\langle \eta, \nabla^2 \tilde{H}(x) \eta \right\rangle. \tag{A.20}$$

The first step is to extend $\tilde{H}$ till second order. Therefore, we start with $\nabla \tilde{H}$ and observe by comparing with (A.12)

$$\begin{aligned} \nabla \tilde{H}(x) = \nabla H(x) &+ \partial_s H^+_{\mathrm{loc}}(s,u) \frac{x_{\mathsf{s}}}{s} - \partial_u H^+_{\mathrm{loc}}(s,u) D h_{\mathsf{s}}^\top(x_{\mathsf{s}}) \frac{x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}})}{u} \\ &+ \partial_u H^+_{\mathrm{loc}}(s,u) \frac{x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}})}{u}, \end{aligned} \tag{A.21}$$

The terms on the first line on the right hand in (A.21) are smooth C^1-functions. Therefore, we can write

$$\nabla \tilde{H}(x) = \partial_u H^+_{\mathrm{loc}}(s,u) \frac{x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}})}{u} + R_1(x), \tag{A.22}$$

where $R_1(x) \in C^1(\mathbb{R}^n)$ and $\|R_1\|_{C^1(\mathbb{R}^n)} \leq C\,|x|$ for some constant C depending on H and h_{s} with their first and second derivatives as well as on p and q. Hence, we note that $|\nabla \tilde{H}(x)| = O(1)$. Now, we calculate the Hessian of $\tilde{H}$ by only considering the leading term of (A.22)

$$\begin{aligned} \nabla^2 \tilde{H}(x) = DR_1(x) &+ \partial^2_{s,u} H^+_{\mathrm{loc}}(s,u) \frac{x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}})}{u} \otimes \frac{x_{\mathsf{s}}}{s} \\ &+ \partial^2_u H^+_{\mathrm{loc}}(s,u) \frac{x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}})}{u} \otimes \frac{(\Pi_{\mathsf{u}} - D h_{\mathsf{s}}^\top(x_{\mathsf{s}}))(x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}}))}{u} \\ &+ \frac{\partial_u H^+_{\mathrm{loc}}(s,u)}{u} \left(\Pi_{\mathsf{u}} - D h_{\mathsf{s}}^\top(x_{\mathsf{s}}) - \frac{x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}})}{u} \otimes \frac{(\Pi_{\mathsf{u}} - D h_{\mathsf{s}}^\top)(x_{\mathsf{s}})(x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}}))}{u} \right). \end{aligned} \tag{A.23}$$

The terms in the first two lines of (A.23) are all at least $O(1)$ by the assumptions (A_p), (B_p), (A_q) and (B_q) as well as by the properties of h_{s} given in its Definition (A.8). The prefactor $\frac{\partial_u H^+_{\mathrm{loc}}(s,u)}{u}$ of the last term in (A.23) blows up like $-\frac{1}{u}$, since $\partial_u H^+_{\mathrm{loc}}(s,u) < 0$ by assumption (A_q). The matrix in the last line is of the form $A - v \otimes Av$,where $A = \mathrm{Id} - D h_{\mathsf{s}}^\top(x_{\mathsf{s}})$ and $v = \frac{x_{\mathsf{u}} - h_{\mathsf{s}}(x_{\mathsf{s}})}{u} \in S^{n-1}$. We can assume that $A > 0$, because of $\left| D h_{\mathsf{s}}^\top(x_{\mathsf{s}}) \right| = O(s)$ by (A.8). For a matrix $A > 0$ holds $A - v \otimes Av \geq 0$ with equality only if v is an eigenvector of A. If we use these observations together with (A.20), we arrive at

$$-C - \left| \frac{\partial_u H^+_{\mathrm{loc}}(s,u)}{u} \right| \leq \tilde{\lambda}_1(x) \leq \tilde{\lambda}_n(x) \leq C,$$

where C is constant depending on $|DR_1(x)|$, $H^+_{\mathrm{loc}}(s,u)$ with its first and second derivatives and h_s with its first derivative. Now, we estimate with the definition of H^+_{loc} and with (B_q)

$$\partial_u H^+_{\mathrm{loc}}(s,u) = -a\delta\, p\left(\frac{s}{\delta}\right)\left|q'\left(\frac{u}{\delta}\right)\right| \geq -2a\delta\, p\left(\frac{s}{\delta}\right).$$

In combination with (A.19), we arrive at a pointwise version of ($\mathbf{\tilde{H}4}$)

$$-(n-1)\left(C + \frac{2a\delta p\left(\frac{s}{\delta}\right)}{u}\right) \leq \nabla\cdot\frac{\nabla\tilde{H}(x)}{|\nabla\tilde{H}(x)|} \leq (n-1)\,C.$$

In the last step of the proof of ($\mathbf{\tilde{H}4}$), we have to combine the estimate A.2 with the trajectories $\psi_\eta(t)$ in $\mathcal{U}$. Therefore, we introduce the functions $s(t)$ and $u(t)$ given by

$$s(t) := |\Pi_{\mathsf{s}}\psi_\eta(t)| \qquad \text{and} \quad u(t) := |\Pi_{\mathsf{u}}\psi_\eta(t) - h_{\mathsf{s}}\left(\Pi_{\mathsf{s}}\psi_\eta(t)\right)|\,.$$

Further we denote with $p_\delta(\cdot) = p(\frac{\cdot}{\delta})$. To finish the proof of ($\mathbf{\tilde{H}4}$), we have to show that for a trajectory $\psi_\eta(t)$ in $\mathcal{U}$ holds

$$\frac{p_\delta(s(t))}{u(t)} = O\left(\frac{1}{(T_\eta - t)^2}\right). \tag{A.24}$$

Now, to show (A.24), we deduce first two differential inequalities for $u(t)$ and $s(t)$ which holds for δ sufficiently small and a satisfying (A.17)

$$1 \geq \dot{s}(t) \geq \frac{\lambda^+_{\mathrm{min}}}{2}s + O(\delta^2), \tag{A.25}$$

$$1 \geq -\dot{u}(t) \geq a(\delta - 2u(t))_+ p_\delta(s(t)) + O(\delta). \tag{A.26}$$

The deduction of (A.26) and (A.25) is similar to the estimate of N_{s} and N_{u}. By direct calculation and term-wise estimation of $|\nabla\tilde{H}|$ in (A.21), we find that $|\nabla\tilde{H}| = O(\delta)$ in $\mathcal{U}$. Hence we can assume from now that $|\nabla\tilde{H}| \leq 1$. The starting point for all estimates is the representation of $\nabla\tilde{H}$ given in (A.11)

$$\begin{aligned}
\dot{s}(t) &= \frac{1}{s(t)}\left\langle \Pi_{\mathsf{s}}\psi_\eta(t), \dot{\psi}_\eta(t)\right\rangle = \frac{1}{s(t)|\nabla\tilde{H}(\psi_\eta(t))|}\left\langle \Pi_{\mathsf{s}}\psi_\eta(t), \nabla\tilde{H}(\psi_\eta(t))\right\rangle \\
&= \frac{1}{s(t)|\nabla\tilde{H}(\psi_\eta(t))|}\left(\langle \Pi_{\mathsf{s}}\psi_\eta(t), N_{\mathsf{s}}(\psi_\eta(t))\rangle + \langle \Pi_{\mathsf{s}}\psi_\eta(t), R_2(x)\rangle\right) \\
&\geq \frac{1}{s(t)}\langle \Pi_{\mathsf{s}}\psi_\eta(t), N_{\mathsf{s}}(\psi_\eta(t))\rangle + O(\delta^2).
\end{aligned}$$

We follow along the lines of (A.14) and (A.17) to estimate $\langle\psi_\eta(t), N_s(\psi_\eta(t))\rangle$ and find

$$\begin{aligned}
\left\langle\frac{\Pi_{\mathsf{s}}\psi_\eta(t)}{s(t)}, N_s(\psi_\eta(t))\right\rangle &= \left\langle\frac{\Pi_{\mathsf{s}}\psi_\eta(t)}{s(t)}, \nabla^2 H(0)\psi_\eta(t)\right\rangle + \partial_s H^+_{\mathrm{loc}}(s(t),u(t))\frac{|\Pi_{\mathsf{s}}\psi_\eta(t)|^2}{s(t)^2} \\
&\quad - \partial_u H^+_{\mathrm{loc}}(s,u)\left\langle\frac{\Pi_{\mathsf{s}}\psi_\eta(t)}{s(t)}, Dh_{\mathsf{s}}^\top(\Pi_{\mathsf{s}}\psi_\eta(t))\frac{\Pi_{\mathsf{u}}\psi_\eta(t) - h_{\mathsf{s}}(\Pi_{\mathsf{s}}\psi_\eta(t))}{u(t)}\right\rangle \\
&\overset{(A.14)}{\geq} \lambda^+_{\mathrm{min}}s(t) - a\delta\left|p'\left(\frac{s(t)}{\delta}\right)\right| q\left(\frac{u(t)}{\delta}\right) - a\delta p\left(\frac{s(t)}{\delta}\right)q'\left(\frac{u(t)}{\delta}\right)\left|Dh_{\mathsf{s}}^\top(\Pi_{\mathsf{s}}\psi_\eta(t))\right| \\
&\overset{(A.15),(A.16)}{\geq} \lambda^+_{\mathrm{min}}s(t) - 8as(t) - 2aC_h s(t).
\end{aligned}$$

Using the same choice of a as in (A.17) shows (A.25). For the differential inequality (A.26) we have to take care of the sign of $\dot{u}(t)$

$$
\begin{aligned}
-\dot{u}(t) &= -\left\langle \frac{\Pi_{\mathbf{u}}\psi_\eta(t) - h_s(\Pi_{\mathbf{s}}\psi_\eta(t))}{u(t)}, \Pi_{\mathbf{u}}\dot{\psi}_\eta(t) - Dh_{\mathbf{s}}(\Pi_{\mathbf{s}}\psi_\eta(t))\,\Pi_{\mathbf{s}}\dot{\psi}_\eta(t) \right\rangle \\
&\geq -\left\langle \frac{\Pi_{\mathbf{u}}\psi_\eta(t) - h_s(\Pi_{\mathbf{s}}\psi_\eta(t))}{u(t)}, \frac{N_{\mathbf{u}}}{|\nabla\tilde{H}(\psi_\eta(t))|} \right\rangle + |Dh_{\mathbf{s}}(\Pi_{\mathbf{s}}\psi_\eta(t))| + O(\delta^2).
\end{aligned}
$$

Since $|Dh_{\mathbf{s}}(\Pi_{\mathbf{s}}\psi_\eta(t))| = O(|\Pi_{\mathbf{s}}\psi_\eta(t)|) = O(\delta)$ by (A.8), we only have to consider the first term. We can use a similar estimate to (A.18) and find

$$
\begin{aligned}
-\left\langle \frac{\Pi_{\mathbf{u}}\psi_\eta(t) - h_s(\Pi_{\mathbf{s}}\psi_\eta(t))}{u(t)}, \frac{N_{\mathbf{u}}}{|\nabla\tilde{H}(\psi_\eta(t))|} \right\rangle &\geq |\lambda^-_{\min}|u(t) + a\delta p\left(\frac{s(t)}{\delta}\right)\left|q'\left(\frac{s(t)}{\delta}\right)\right| \\
&\overset{(\mathrm{B}_q)}{\geq} |\lambda^-_{\min}|u(t) + a\delta\left(1 - 2\frac{u(t)}{\delta}\right)_+ p\left(\frac{s(t)}{\delta}\right),
\end{aligned}
$$

which concludes the proof of (A.26).

Let us introduce the entrance time E_η of $\psi_\eta(t)$ into $\operatorname{supp} H^+_{\mathrm{loc}}$ where we can assume that $\operatorname{supp} H^+_{\mathrm{loc}} \subset B_{2\delta}$. From (A.25) follows that the function $s(t)$ is strictly increasing, whenever $s(t)$ is large enough and δ small enough. Let us assume that δ is chosen small enough such that $\dot{s}(t) \geq c_s > 0$ whenever $s(t) \geq \frac{\delta}{2}$. Then, we can consider two cases: First, there is $t_0 \in (E_\eta, T_\eta)$ with $s(t_0) \geq \frac{\delta}{2}$ and hence $s(t)$ is increasing for $t \in (t_0, T_\eta)$ and second there is no such t_0 and $s(t) \in (0, \frac{\delta}{2})$ for all $t \in (E_\eta, T_\eta)$. Let us first consider the latter case. Further, as we are only interested in the blow-up behavior, we can assume w.l.o.g. $u(t) \leq \frac{\delta}{4}$ and find

$$
-\dot{u}(t) \geq \frac{a\delta}{2} p\left(\frac{1}{2}\right) + O(\delta^2) \geq c_u > 0. \tag{A.27}
$$

Hence, especially $u(t) \geq c_u(T_\eta - t)$ from which we deduce with (A_p)

$$
\frac{p_\delta(s(t))}{u(t)} \leq \frac{1}{c_u(T_\eta - t)}
$$

and therewith also (A.24). Now, we assume that there exists $t_0 \in (E_\eta, T_\eta)$ with $s(t_0) > \frac{\delta}{2}$. We use $u(t) \geq \frac{a\delta}{2}\int_t^{T_\eta} p(\frac{s(t')}{\delta})\mathrm{d}t'$ for $t \in (t_0, T_\eta)$ by (A.26). Then it holds

$$
\frac{p_\delta(s(t))}{u(t)} \leq \frac{2}{a\delta}\frac{p_\delta(s(t))}{\int_t^{T_\eta} p_\delta(s(t'))\mathrm{d}t'} \overset{(\mathrm{A.25})}{\leq} \frac{2}{a\delta}\frac{p_\delta(s(t))}{\int_{s(t)}^{s(T_\eta)} p_\delta(s)\mathrm{d}s} \overset{(\mathrm{D}_p)}{\leq} \frac{2C_p}{a\delta}\frac{1}{(s(T_\eta) - s(t))^2}.
$$

By using the lower bound $\dot{s}(t) \geq c_s > 0$, we can infer

$$
s(T_\eta) - s(t) = \int_t^{T_\eta} \dot{s}(\tau)\mathrm{d}\tau \geq c_s(T_\eta - t),
$$

from which the estimate (A.24) follows.

Argument for (**$\tilde{\mathbf{H}}$5**): We conclude in several steps. Firstly, we deduce a pointwise estimate

$$\langle \nabla H(x), \nabla \tilde{H}(x)\rangle \geq \frac{(\lambda^+_{\min})^2}{2} s^2 + |\lambda^-_{\min}| a\delta \left(1 - \frac{2u}{\delta}\right)_+ p_\delta(s) u + \left(\lambda^-_{\min}\right)^2 u^2 + O(s^2\delta) + O(u^2\delta). \tag{A.28}$$

Then, in the second step, we get rid of the error terms, as they can be compensated by the first and second term of (A.28), respectively. If we for the moment assume that (A.28) holds, we still need to recombine this estimate (A.28) with the trajectories. Hereby, we use again (A.25) and (A.26). Like in deducing (**$\tilde{\mathbf{H}}$4**), we have to consider two cases: First, there exists $t_0 \in (E_\eta, T_\eta)$ with $s(t_0) \geq \frac{1}{2}\delta$ and secondly, there exists no such t_0. In the first case, there is nothing to show, since we can assume that $s(t)$ is monotone and therefore with (A.28) holds

$$\forall t \in (t_0, T_\eta) : \langle \nabla H(\psi_\eta(t)), \nabla \tilde{H}(\psi_\eta(t))\rangle \geq \frac{(\lambda^+_{\min})^2}{4} s(t_0)^2 \geq \frac{(\lambda^+_{\min})^2}{16}\delta^2.$$

Now, we can assume, that $s(t) < \frac{1}{2}\delta$ for all $t \in (E_\eta, T_\eta)$ or for all $t \in (E_\eta, t_0)$. Therefore it holds (A.27) and we can deduce the estimate for $u \leq \frac{\delta}{4}$

$$\langle \nabla H(\psi_\eta(t)), \nabla \tilde{H}(\psi_\eta(t))\rangle \geq |\lambda^-_{\min}| a \frac{\delta}{2} p\left(\frac{1}{2}\right) u(t) \geq |\lambda^-_{\min}| a \frac{\delta}{2} p\left(\frac{1}{2}\right) c_u (T_\eta - t).$$

The conclusion (**$\tilde{\mathbf{H}}$5**) follows from the observation that $|\nabla \tilde{H}| = O(\delta)$ in $\mathcal{U}$ and therefore especially $|\nabla \tilde{H}| \leq 1$ for δ chosen appropriate small enough.

To finish the proof, we still have to show (A.28). We start with (A.11) and (A.12) to deduce

$$\begin{aligned}\left\langle \nabla H(x), \nabla \tilde{H}(x)\right\rangle &= \left\langle \nabla^2 H(0)(x_{\mathtt{s}} + x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})) + R_2(x), N_{\mathtt{s}}(x) + N_{\mathtt{u}}(x) + R_2(x)\right\rangle \\ &= \left\langle \nabla^2 H(0)(x_{\mathtt{s}} + x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})), N_{\mathtt{s}}(x) + N_{\mathtt{u}}(x)\right\rangle \\ &\quad + \left\langle R_2(x), N_s(x) + N_u(x) + \nabla^2 H(0)(x_{\mathtt{s}} + x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})) + R_2(x)\right\rangle.\end{aligned}$$

Let us first estimate the order of the error terms containing $R_2(x)$. With (A.11) we find that $|R_2(x)| = O(|x|^2) = O(s^2) + O(u^2)$. Further, we have $|N_s(x)| = O(\delta)$ as well as $|N_u(x)| = O(\delta)$ and of course $|x| = O(\delta)$. Therefore, the total approximation error can be estimated by

$$\left\langle R_2(x), N_s(x) + N_u(x) + \nabla^2 H(0)(x_{\mathtt{s}} + x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})\right\rangle = O(\delta s^2) + O(u^2\delta).$$

We find an estimate for $\left\langle \nabla^2 H(0)(x_{\mathtt{s}} + x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})), N_{\mathtt{s}}(x) + N_{\mathtt{u}}(x)\right\rangle$by using the representation of $\nabla\tilde{H}$ in (A.12) and going along the lines of (A.14) and (A.18)

$$\begin{aligned}
&\left\langle \nabla^2 H(0)(x_{\mathtt{s}} + x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})), N_{\mathtt{s}}(x) + N_{\mathtt{u}}(x)\right\rangle \\
&= \left\langle \nabla^2 H(0) x_{\mathtt{s}}, N_{\mathtt{s}}\right\rangle + \left\langle \nabla^2 H(0)\left(x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})\right), N_{\mathtt{u}}(x)\right\rangle \\
&\geq \left\langle \nabla^2 H(0) x_{\mathtt{s}}, \nabla^2 H(0) x_{\mathtt{s}}\right\rangle \\
&\qquad\qquad - a\delta \left|p'\left(\frac{s}{\delta}\right)\right| q\left(\frac{u}{\delta}\right) \left\langle \nabla^2 H(0) x_{\mathtt{s}}, \frac{x_{\mathtt{s}}}{s}\right\rangle \\
&\qquad\qquad - a\delta p\left(\frac{s}{\delta}\right) \left|q'\left(\frac{u}{\delta}\right)\right| \left|\nabla^2 H(0) x_{\mathtt{s}}\right| \left|Dh_{\mathtt{s}}^{\top}(x_{\mathtt{s}})\right| \\
&\qquad + \left\langle \nabla^2 H(0)\left(x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})\right), \nabla^2 H(0)\left(x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})\right)\right\rangle \\
&\qquad\qquad + a\delta p\left(\frac{s}{\delta}\right) \left|q'\left(\frac{u}{\delta}\right)\right| \left\langle -\nabla^2 H(0)\left(x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})\right), \frac{x_{\mathtt{u}} - h_{\mathtt{s}}(x_{\mathtt{s}})}{u}\right\rangle \\
&\geq (\lambda^+_{\min})^2 s^2 - 8a\lambda^+_{\max} s^2 - 2a\delta\lambda^+_{\max} C_h s^2 + (\lambda^-_{\min})^2 u^2 + a\delta p\left(\frac{s}{\delta}\right)\left(1 - \frac{2u}{\delta}\right)_+ \left|\lambda^-_{\min}\right| u.
\end{aligned}$$

We finish by estimating the first three terms. Therefore, we use, like in (A.17), the assumption that $\delta \leq C_h^{-1}$ and obtain

$$(\lambda^+_{\min})^2 s^2 - 8a\lambda^+_{\max} s^2 - 2a\lambda^+_{\max}\delta C_h s^2 \geq \left((\lambda^+_{\min})^2 - 10a\lambda^+_{\max}\right) s^2 \geq \frac{(\lambda^+_{\min})^2}{2} s^2,$$

where we set $a \leq \frac{(\lambda^+_{\min})^2}{20\lambda^+_{\max}}$ in the last estimate, which leads to (A.28) and finishes the proof. □

B. Gaussian integrals and linear algebra

B.1. Partial Gaussian integrals

This section is devoted to proof the representation for partial or incomplete Gaussian integrals. The formula (B.1) is an ingredient to evaluate the weighted transport cost in Section 3.3.

Lemma B.1 (Partial Gaussian integral)**.** *Let $\Sigma^{-1} \in S^+(n)$ be a symmetric positive definite matrix and let $\eta \in S^{n-1}$ be a unit vector. Therewith, $\left\{r\eta + z^{\perp}\right\}_{r\in\mathbb{R}}$ is for $z^{\perp} \in \operatorname{span}\{\eta\}^{\perp}$ an affine subspace of $\mathbb{R}^n$. The integral of a centered Gaussian w.r.t. to this subspace is given by*

$$\int_{\mathbb{R}} \exp\left(-\frac{1}{2}\Sigma^{-1}[r\eta + z^{\perp}]\right) \mathrm{d}r = \frac{\sqrt{2\pi}}{\sqrt{\Sigma^{-1}[\eta]}} \exp\left(-\tilde{\Sigma}^{-1}[z^{\perp}]\right),$$
$$\text{with} \quad \tilde{\Sigma}^{-1} = \Sigma^{-1} - \frac{\Sigma^{-1}\eta \otimes \Sigma^{-1}\eta}{\Sigma^{-1}[\eta]}.$$

Proof. To evaluate this integral on an one-dimensional subspace of $\mathbb{R}^n$, we have to expand the quadratic form $\Sigma^{-1}[r\eta + z^\perp]$ and arrive at the relation

$$\begin{aligned}
&\int_{\mathbb{R}} \exp\left(-\frac{1}{2}\Sigma^{-1}[r\eta + z^\perp]\right) \mathrm{d}r \\
&\quad = \exp\left(-\frac{1}{2}\Sigma^{-1}[z^\perp]\right) \int_{\mathbb{R}} \exp\left(-\frac{r^2}{2}\Sigma^{-1}[\eta] + r\left\langle \eta, \Sigma^{-1}z^\perp\right\rangle\right) \mathrm{d}r \\
&\quad = \exp\left(-\frac{1}{2}\Sigma^{-1}[z^\perp]\right) \frac{\sqrt{2\pi}}{\sqrt{\Sigma^{-1}[\eta]}} \exp\left(\frac{\left\langle \eta, \Sigma^{-1}z^\perp\right\rangle^2}{2\Sigma^{-1}[\eta]}\right) \\
&\quad = \frac{\sqrt{2\pi}}{\sqrt{\Sigma^{-1}[\eta]}} \exp\left(-\frac{1}{2}\left(\Sigma^{-1} - \frac{\Sigma^{-1}\eta \otimes \Sigma^{-1}\eta}{\Sigma^{-1}[\eta]}\right)[z^\perp]\right),
\end{aligned}$$

which concludes the hypothesis. $\square$

B.2. Subdeterminants, adjugates and inverses

Let $A \in S^+(n)$, then define for $\eta \in S^{n-1}$ the matrix

$$\tilde{A} := A - \frac{A\eta \otimes A\eta}{A[\eta]}. \tag{B.1}$$

The matrix $\tilde{A}$ has at least rank $n-1$, since we subtracted from the positive definite matrix A a rank-1 matrix. Further, from the representation it is immediate, that $\tilde{A}$ has rank $n-1$ if and only if η is an eigenvector of A. In this case $\ker A = \operatorname{span}\eta$. It is easy to show that

$$\tilde{A} > 0 \quad \text{on } \operatorname{span}\{\eta\}^\perp.$$

Let $V = \operatorname{span}\{\eta\}^\perp$ be the $(n-1)$-dimensional subspace perpendicular to η. Then for a matrix $A \in S^+(n)$ we want to calculate the determinant of A restricted to this subspace V. This determinant is obtained by first choosing $Q \in SO^n$ such that $Q(\{0\}\times\mathbb{R}^{n-1}) = V$ and then evaluating the determinant of the minor consisting of the $(n-1)\times(n-1)$ lower right submatrix of $Q^\top AQ$ denoted by $\det_{1,1}(Q^\top AQ)$. Hence, we have

$$\det\nolimits_{1,1}(Q^\top AQ), \qquad \text{with } Q \in SO(n) : Q^\top\eta = e^1 = (1,0,\dots,0)^\top.$$

Since $V = \operatorname{span}\{\eta\}^\perp$, it follows that the first column of Q is given by η and we can decompose $Q^\top AQ$ into

$$Q^\top AQ = \begin{pmatrix} A[\eta] & \widehat{Q^\top A\eta} \\ \widehat{Q^\top A\eta}^\top & \widehat{Q^\top AQ} \end{pmatrix},$$

where for a matrix M, $\widehat{M}$ is the lower right $(n-1)\times(n-1)$ submatrix of M and for a vector v, $\widehat{v}$ the $(n-1)$ lower subvector of v. Therewith, we find a similarity transformation which applied to $Q^\top AQ$ results in

$$\begin{aligned}\det A = \det Q^\top AQ &= \det\left(\begin{pmatrix} A[\eta] & \widehat{Q^\top A\eta} \\ \widehat{Q^\top A\eta}^\top & \widehat{Q^\top AQ}\end{pmatrix}\begin{pmatrix} 1 & -\frac{\widehat{Q^\top A\eta}}{A[\eta]} \\ 0 & \mathrm{Id}_{n-1}\end{pmatrix}\right) \\ &= \det\begin{pmatrix} A[\eta] & 0 \\ \widehat{Q^\top A\eta}^\top & \widehat{Q^\top AQ} - \frac{\widehat{A\eta}\otimes\widehat{A\eta}}{A[\eta]}\end{pmatrix} = A[\eta]\det{}_{1,1}\left(Q^\top AQ - \frac{Q^\top A\eta\otimes Q^\top A\eta}{A[\eta]}\right).\end{aligned}$$

The determinant of the minor is given by

$$\det{}_{1,1}\left(Q^\top AQ - \frac{Q^\top A\eta\otimes Q^\top A\eta}{A[\eta]}\right) = \det{}_{1,1}\left(Q^\top\left(A - \frac{A\eta\otimes A\eta}{A[\eta]}\right)Q\right).$$

Hence, by the definition (B.1) of $\tilde{A}$ and the subdeterminant we found the identity

$$\det A = A[\eta]\det{}_{1,1}(Q^\top AQ). \tag{B.2}$$

B.3. A matrix optimization

Lemma B.2. *Let $B \in S^+(n-1)$, then it holds*

$$\inf_{A\in S^+(n-1)}\left\{\frac{\det A}{\sqrt{\det(2A-B)}} : 2A > B\right\} = \sqrt{\det B}$$

and for the optimal A holds $A = B$.

Proof. We note that

$$\frac{\det A}{\sqrt{\det(2A-B)}} = \frac{1}{\sqrt{\det(A^{-1})\det(2\,\mathrm{Id} - A^{-\frac{1}{2}}BA^{-\frac{1}{2}})}}.$$

Therewith, it is enough to maximize the radical of the root. Therefore, we substitute $A^{-\frac{1}{2}} = CB^{-\frac{1}{2}}$ with $C > 0$ not necessarily symmetric and observe that $A^{-\frac{1}{2}} = B^{-\frac{1}{2}}C^\top$. We obtain

$$\det(A^{-1})\det(2\,\mathrm{Id} - A^{-\frac{1}{2}}BA^{-\frac{1}{2}}) = \det(B^{-1})\det(CC^\top)\det(2\,\mathrm{Id} - CC^\top).$$

Note, that $CC^\top \in S^+(n)$ and it is enough to calculate

$$\sup_{\tilde{C}\in S^+(n)}\left\{\det(\tilde{C})\det(2\,\mathrm{Id} - \tilde{C}) : \tilde{C} < 2\,\mathrm{Id}\right\}.$$

From the constraint $0 < \tilde{C} < 2\,\mathrm{Id}$ we can write $\tilde{C} = \mathrm{Id} + D$, where D is symmetric and satisfies $-\mathrm{Id} < D < \mathrm{Id}$ in the sense of quadratic forms. From here, we finally observe

$$\det(\tilde{C})\det(2\,\mathrm{Id} - \tilde{C}) = \det(\mathrm{Id} + D)\det(\mathrm{Id} - D) = \det(\mathrm{Id} - D^2).$$

Since $D^2 \geq 0$, we find the optimal $\tilde{C}$ given by Id, which yields that $A = B$. $\square$

B.4. Jacobi's formula

Lemma B.3 (Jacobi's formula)**.** *Let* $\mathbb{R} \ni t \mapsto \Phi_t \in \{A \in \mathbb{R}^{n\times n} : \det A \neq 0\}$ *be a differentiable function, then*

$$\frac{\mathrm{d}}{\mathrm{d}t} \ln \det \Phi_t = \operatorname{tr}\left(\Phi_t^{-1}\dot{\Phi}_t\right). \tag{B.3}$$

Proof. We first note that the determinant of $\Phi(t)$ is a multilinear function d of the columns $\phi_t^1, \dots, \phi_t^n$, i.e. $\det \Phi_t = d(\phi_t^1, \dots, \phi_t^n)$ Then, it follows

$$\frac{\mathrm{d}}{\mathrm{d}t} \det \Phi_t = d(\dot{\phi}_t^1, \phi_t^2, \dots, \phi_t^n) + \dots + d(\phi_t^1, \dots, \phi_t^{n-1}, \dot{\phi}_t^n).$$

Now, the proof consists of two steps. We first proof the identity (B.3) for $\Phi_t = \mathrm{Id}$ and then generalize this result. If we assume w.l.o.g. that $\Phi_0 = \mathrm{Id}$. By expanding the determinant $d(\dot{\phi}_t^1, \phi_t^2, \dots, \phi_t^n)$ along its first column it immediately follows that

$$d(\dot{\phi}_t^1, \phi_t^2, \dots, \phi_t^n) = \dot{\phi}_t^{1,1}.$$

From here we conclude that

$$\frac{\mathrm{d}}{\mathrm{d}t} \det \Phi_t = \operatorname{tr} \dot{\Phi}_t.$$

Now, let $\Phi_t = A$ be a general invertible matrix. Hence, we can apply the result from the first step to $A^{-1}\Phi_t$ and arrive at

$$\frac{\mathrm{d}}{\mathrm{d}t} \det\left(A^{-1}\Phi_t\right) = \operatorname{tr}\left(A^{-1}\dot{\Phi}_t\right).$$

The results follows by substituting A back. ☐

C. Existence of a spectral gap for L

In this short part of the appendix, we state a standard argument that ensures that the conditions (**A1**) and (**A2**) imply a spectral gap. More precisely, we make the following definition of the spectral gap

Definition C.1. We say that the operator $L = \varepsilon\Delta - \nabla H \cdot \nabla$ given by (1.1) satisfies a spectral gap of order $\varrho > 0$ i.e. $\mathrm{SG}(\varrho)$, if L has discrete non-negative spectrum and the order eigenvalues λ_i satisfy

$$\lambda_0 = 0 < \lambda_1 \leq \lambda_2 \leq \dots \qquad \text{and} \qquad \lambda_1 \geq \varrho.$$

We take over the argument of [Kun02, Proposition 3.7] and adapt it to the case of small noise ε.

Proposition C.2. *Assume that H satisfies (**A1**) and (**A2**) with constants $C_H, K_H > 0$. Then for all $\varepsilon \leq \frac{1}{2}\frac{C_H^2}{2c+K_H+C_H^2}$ with $c > 0$ the operator $L = \varepsilon\Delta - \nabla H \cdot \nabla$ given by (1.1) satisfies* $\mathrm{SG}(\varrho)$ *for some $\varrho > 0$.*

Proof. The operator $L = \varepsilon\Delta - \nabla H \cdot \nabla$ on $L^2(\mu)$ can be transformed into a *Schrödinger operator* with potential

$$L_S := -\varepsilon\Delta + \tfrac{1}{4\varepsilon}\left|\nabla H\right|^2 - \tfrac{1}{2}\Delta H$$

on $L^2(\mathrm{d}x)$. This can be seen by using the unitary transformation $U : L^2(\mathrm{d}x) \to L^2(\mu)$ given by $f \mapsto \exp(\frac{H}{2\varepsilon})f$. So, by partial integration we get

$$\begin{aligned}\mathcal{E}(Uf, Ug) &= \varepsilon \int \nabla f \cdot \nabla g + \tfrac{1}{4\varepsilon^2}|\nabla H|^2 fg + \tfrac{1}{2\varepsilon}\nabla H \cdot (f\nabla g + g\nabla f)\,\mathrm{d}x \\ &= \int \varepsilon\nabla f \cdot \nabla g + \left(\tfrac{1}{4\varepsilon}|\nabla H|^2 - \tfrac{1}{2}\Delta H\right) fg\,\mathrm{d}x = \int (L_S f)g\,\mathrm{d}x.\end{aligned}$$

For the Schrödinger operator L_S it is known (see e.g. [BS91, Theorem 3.1]) that

$$\liminf_{|x|\to\infty} \left(\tfrac{1}{4\varepsilon}|\nabla H|^2 - \tfrac{1}{2}\Delta H\right) \geq c > 0 \tag{C.1}$$

is a sufficient condition to have a discrete spectrum on $(-\infty, c)$ and in addition for every $c' < c$ and $C' < \infty$ to have a finite spectrum on $(-C', c')$. The condition (C.1) is implied by the assumption (**A2**) for $\varepsilon \leq \frac{1}{2}\frac{C_H^2}{2c+K_H+C_H^2}$ (see the proof of Lemma 4.6 for a similar argument). Since the transformation U was unitary L_S and $-L$ have the same spectrum. Hence, L has also a discrete spectrum on $(-c, \infty)$ and as it is a non-negative operator it has a discrete spectrum on $[0, \infty)$ which is finite on $[0, C')$ for every $C' < \infty$, which implies a spectral gap. □

D. Conventions and notation

$\mathrm{d}x$	n-dimensional Lebesgue measure
$\mathcal{H}^k(\mathrm{d}x)$	k-dimensional Hausdorff measure
$\mu(\cdot)$	density of of the probability measure μ
$\mu(\mathrm{d}x)$	short form for $\mu(\cdot)\mathrm{d}x$
$\mu(\cdot\vert y)$	conditional measure obtained from μ by conditioning on y
Z	a generic partition sum normalizing a probability measure
$\operatorname{supp}\mu$	the support of μ, $\operatorname{supp}\mu = \overline{\{\mu(x) > 0\}}$
$\mathbb{1}_\Omega$	characteristic function on Ω
$\mu\llcorner\Omega$	restriction of μ onto $\Omega \subset \operatorname{supp}\mu$, $(\mu\llcorner\Omega)(\mathrm{d}x) := \frac{\mathbb{1}_\Omega(x)}{\mu(\Omega)}\mu(x)\mathrm{d}x$
$\Phi_\sharp\mu$	push-forward measure under $\Phi : \mathbb{R}^n \to \mathbb{R}^n \supset U$, $(\Phi_\sharp\mu)(U) = \mu(\Phi^{-1}(U))$
$A^\top$	matrix transpose
$\operatorname{tr} A$	trace of A
$\langle x, y\rangle = x \cdot y$	Euclidean scalar product between $x, y \in \mathbb{R}^n$
$\lvert\cdot\rvert$	Euclidean norm
$A[x]$	testing a quadratic form $A[x] = \langle x, Ax\rangle$
$\det_{i,j} A$	determinant of the matrix A with i-th row and j-th column removed
$A \leq B$	$A[x] \leq B[x]$ for all $x \in \mathbb{R}^n$
$S^+(n)$	set of symmetric positive definite matrices, i.e. $A = A^\top$ and $A > 0$.
S^{n-1}	unit sphere in $\mathbb{R}^n$, i.e. $\{\eta \in \mathbb{R}^n : \lvert x\rvert = 1\}$
$\sphericalangle(\eta, \theta)$	geodesic on S^{n-1} between $\eta, \theta \in S^{n-1}$
$SO(n)$	rotational matrices in $\mathbb{R}^n$, $Q \in \mathbb{R}^{n\times n}$: $Q^{-1} = Q^\top$, $\det Q = 1$.
$\nabla^2 H$	Hessian of $H : \mathbb{R}^n \to \mathbb{R}$
$\nabla \cdot F$	the divergence of $F : \mathbb{R}^n \to \mathbb{R}^n$
$\dot{\psi}_t$	derivative in parameter $t \in \mathbb{R}$
$\omega(\varepsilon)$	a smooth monotone decreasing function satisfying $\geq \lvert\ln\varepsilon\rvert^{\frac{1}{2}}$ for $\varepsilon < 1$
$\approx, \lesssim, \gtrsim$	$=, \leq, \geq$ up to multiplicative error of the form $1 + O(\sqrt{\varepsilon}\,\omega^3(\varepsilon))$.
C, c	constants only depending on the dimension n and H
C_f, c_f	constants additionally depending on the function or variable f

Acknowledgment

This work is part of the Ph.D. thesis of the second author, written under the supervision of Stephan Luckhaus at the University of Leipzig.
The authors are greatly indebted to Felix Otto for drawing the authors' attention to the low temperature regime and several helpful discussions.
They wish to thank the Max-Planck-Institute for Mathematics in the Sciences in Leipzig where the paper was written under great working conditions. The second author thanks the DFG project FOR718, the IMPRS of the Max-Planck-Institute and the University of Leipzig for financial support.

References

[Arr89] S. Arrhenius. *Z. Phys. Chem.*, 4:226, 1889.

[BBCG08] D. Bakry, F. Barthe, P. Cattiaux, and A. Guillin. A simple proof of the Poincaré inequality for a large class of probability measures including the log-concave case. *Electron. Commun. Probab.*, 13:60–66, 2008.

[BCG08] D. Bakry, P. Cattiaux, and A. Guillin. Rate of convergence for ergodic continuous Markov processes: Lyapunov versus Poincaré. *J. Funct. Anal.*, 254(3):727–759, 2008.

[BÉ85] D. Bakry and M. Émery. Diffusions hypercontractives. In *Séminaire de probabilités, XIX, 1983/84*, volume 1123 of *Lecture Notes in Math.*, pages 177–206. Springer, Berlin, 1985.

[BEGK04] A. Bovier, M. Eckhoff, V. Gayrard, and M. Klein. Metastability in reversible diffusion processes. I. Sharp asymptotics for capacities and exit times. *J. Eur. Math. Soc. (JEMS)*, 6(4):399–424, 2004.

[Ber11] N. Berglund. Kramers' law: Validity, derivations and generalisations. *arXiv:1106.5799v1*, 2011.

[BG99] S. G. Bobkov and F. Götze. Exponential integrability and transportation cost related to logarithmic Sobolev inequalities. *J. Funct. Anal.*, 163(1):1–28, 1999.

[BGK05] A. Bovier, V. Gayrard, and M. Klein. Metastability in reversible diffusion processes. II. Precise asymptotics for small eigenvalues. *J. Eur. Math. Soc. (JEMS)*, 7(1):69–99, 2005.

[BS91] F. A. Berezin and M. A. Shubin. *The Schrödinger equation*, volume 66 of *Mathematics and its Applications (Soviet Series).* Kluwer Academic Publishers Group, Dordrecht, 1991.

[CM10] D. Chafaï and F. Malrieu. On fine properties of mixtures with respect to concentration of measure and Sobolev type inequalities. *Ann. Inst. Henri Poincaré Probab. Stat.*, 46(1):72–96, 2010.

[Eyr35] H. Eyring. The activated complex in chemical reactions. *J. Chem. Phys.*, 3(2):107, 1935.

[Fou05] P. Fougères. Spectral gap for log-concave probability measures on the real line. In *Séminaire de Probabilités XXXVIII*, volume 1857 of *Lecture Notes in Math.*, pages 95–123. Springer, Berlin, 2005.

[FW98] M. I. Freidlin and A. D. Wentzell. *Random perturbations of dynamical systems*, volume 260 of *Grundlehren der Mathematischen Wissenschaften.* Springer-Verlag, New York, second edition, 1998.

[GMSvdM02] O. Gonzalez, J. H. Maddocks, F. Schuricht, and H. von der Mosel. Global curvature and self-contact of nonlinearly elastic curves and rods. *Calc. Var. Partial Differential Equations*, 14(1):29–68, 2002.

[GOV87] A. Galves, E. Olivieri, and M. E. Vares. Metastability for a class of dynamical systems subject to small random perturbations. *Ann. Probab.*, 15(4):1288–1305, 1987.

[GOVW09] N. Grunewald, F. Otto, C. Villani, and M. Westdickenberg. A two-scale approach to logarithmic Sobolev inequalities and the hydrodynamic limit. *Ann. Inst. Henri Poincaré Probab. Stat.*, 45(2):302–351, 2009.

[HKN04] B. Helffer, M. Klein, and F. Nier. Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach. *Mat. Contemp.*, 26:41–85, 2004.

[HN05] B. Helffer and F. Nier. *Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians*, volume 1862 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2005.

[HN06] B. Helffer and F. Nier. Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach: the case with boundary. *Mém. Soc. Math. Fr. (N.S.)*, (105):vi+89, 2006.

[HS87] R. Holley and D. Stroock. Logarithmic Sobolev inequalities and stochastic Ising models. *J. Statist. Phys.*, 46:1159–1194, 1987.

[Kra40] H. A. Kramers. Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica*, 7:284–304, 1940.

[Kun02] A. Kunz. On extremes of multidimensional stationary diffusion processes in euclidean norm. *Preprint*, 2002.

[Mic99] L. Miclo. An example of application of discrete Hardy's inequalities. *Markov Process. Related Fields*, 5(3):319–330, 1999.

[Muc72] B. Muckenhoupt. Hardy's inequality with weights. *Studia Math.*, 44:31–38, 1972.

[Olv97] F. W. J. Olver. *Asymptotics and special functions.* AKP Classics. A K Peters Ltd., Wellesley, MA, 1997.

[Tes12] G. Teschl. *Ordinary differential equations and dynamical systems*, volume XXX of *Graduate Studies in Mathematics.* American Mathematical Society, 2012. to appear.